

# Response of the Diatom *Melosira varians* to Domestic Wastewater Effluents

**Type of the Article:** Original Research Article

## Abstract

This study investigates the physiological and ecological responses of the benthic diatom *Melosira varians* to varying concentrations of domestic wastewater effluents under laboratory conditions. Effluent dilutions of 12.5%, 25%, and 50% were used to assess the species' capacity for nutrient removal, growth, morphological alterations, and pigment composition over a 10-day period. The results demonstrated remarkable phosphate removal efficiencies ( $\geq 97\%$ ) across all treatments and a progressive nitrate removal response, reaching up to 70.59% in the highest effluent concentration. Morphometric analysis revealed significant increases in cell width and length, particularly under moderate (25%) and high (50%) effluent exposures. Chlorophyll analysis indicated a predominance of chlorophyll c and a reduction of chlorophyll b and c in response to nutrient-rich conditions, suggesting a possible shift towards heterotrophic or mixotrophic metabolism. Multivariate analysis highlighted strong associations between cell morphology and biomass, while chlorophyll content showed no direct correlation with biomass accumulation. These findings demonstrate the metabolic flexibility and nutrient uptake potential of *M. varians*, reinforcing its applicability in sustainable wastewater treatment systems.

## Keywords:

Diatoms; *Melosira varians*; Domestic effluents.

## 1. INTRODUCTION

Urban development and expansion, industrial activities, agriculture, livestock farming, etc., generate effluents and wastewater that require proper management. When discharged into natural bodies of water, they generate problems of contamination and/or eutrophication, affecting the natural purification processes of these aquatic ecosystems with loss of biodiversity and destruction of these ecosystems.

Eutrophication is a process of deterioration in the quality of water bodies. It is caused by the enrichment of nutrients, mainly nitrogen and phosphorus,

generating major ecological, health and economic impacts on a regional scale [1].

Diatoms are dynamic microorganisms with rich diversity and detailed membrane design. They are the most dominating phytoplankton with an overall number of around 200,000 species having complex variability in dimensions and shapes [2], [3], [4].

Diatoms' distinctive characteristic compared to the phytoplankton community is their silica cell wall, known as a frustule. The silica shell protects them from negative environmental stress, giving them a competitive advantage [5]. When silicate is not a limiting factor, other

microalgae are unable to compete with diatoms.

Patterns of benthic diatom communities are responsive to the nature of the physical and chemical characteristics of lotic systems [6], [7], [8]. They respond rapidly to degradation of water quality, often changing in both taxonomic composition and biomass where even slight contamination occurs [9], [10], [11]

Additionally, diatoms can take up phosphorus faster than other algae, as well as all forms of nitrogen—such as nitrate, nitrite, urea, and ammonia [12].

Phosphorus uptake by benthic diatoms occurs in rapid and continuous episodes (in less than one hour), leading to an increase in the oxygenated layer of sediments and inhibiting phosphorus release, which contributes to the purification of the water column [13].

In eutrophic lakes, it is not surprising that the most abundant diatoms in the benthic community are species with high nutrient assimilation strategies [14].

*Melosira varians* is very common in rivers and lakes. This species is considered to be tolerant of poor water quality, although it may occur in more pristine sites as well. This diatom is primarily found in benthic habitats, with cells joined in long filamentous chains. Both individual cells and filaments may become entrained in the plankton.

*Melosira* had a potential ability to cause the blooms in winter. In this study, the dominant species (or genera) *Melosira varians* is generally considered to be tolerant of low light and ice cover. Nutrient addition, it was shown, can increase the benthic diatom abundance in temperate rivers [15].

Our objective was to study the ability of the diatom *M. varians* to remove nitrogen and phosphorus from domestic effluents.

## 2. MATERIALS AND METHODS

### 2.1. Sampling and Isolation of Microalgae

Samples were collected during the fall-winter period of 2023. For this purpose, sediment was aspirated from different areas of an urban riverbed using a plastic syringe, until a significant amount of sediment was obtained according to standard collection techniques [16].

The samples were filtered on site using a 35  $\mu\text{m}$  mesh sieve and transported in a cooler. One hundred milliliters of each sample were placed in sterile 500 ml bottles and brought to a final volume of 500 ml using sterilized river water. The bottles were kept under natural prevailing conditions at the site for two days, at a temperature of temperature ( $22 \pm 1$  °C), illumination with a 16:8 photoperiod (2,800 lux). Subsequently, two 100 ml samples were taken from each bottle, placed into sterile 500 ml bottles, and brought to a final volume of 500 ml with f/4 culture medium (50% f/2 medium, Guillard, 1975, and 50% sterilized river water) for seven days.

Cell counts were also performed for each sample using a Neubauer haemocytometer (0.1 mm depth) to compare the quantity of algal cells and other microorganisms. The least-contaminated cultures were selected and filtered, 100 ml from each bottle were cultured and brought to a final volume of 500 ml with f/2 culture medium. The culture was purified through serial dilutions.

For taxonomic analysis of the strains, specific literature was consulted: Bellinger

& Sigeo (2010), Lee (2008), Guiry, M.D. & Guiry, G.M. (2025), *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. Available at: <http://www.algaebase.org>.

The *Melosira varians* inocula were placed in 1-liter photobioreactors containing f/2 medium [17] to obtain sufficient biomass under controlled laboratory conditions: temperature ( $22 \pm 1$  °C), illumination with a 16:8 photoperiod (2,800 lux), and daily provision of carbon dioxide for growth. Once the biomass was obtained, it was stored in a refrigerator and later reactivated to carry out the phycoremediation treatments.


## 2.2. Wastewater Sampling

The sampling of wastewater effluents was conducted in accordance with the standardized methods for the Examination of Water and Wastewater [18]. The effluent was collected at the outlet of a local wastewater treatment plant in 5-liter

polypropylene containers. Once collected, samples were refrigerated at 4 °C for subsequent transportation to the laboratory.

The effluent was left to stand in darkness for approximately seven days to induce anaerobic conditions and sedimentation. This process aimed to prevent the proliferation of photoautotrophic organisms and the formation of protozoan cysts and helminth eggs (primary microalgal predators), while simultaneously promoting bacterial concentration [19].

An experimental design was implemented, comprising a control group and increasing concentrations of effluent (12.5%, 25%, and 50%), all inoculated with  $1 \times 10^6$  cells/mL of the obtained biomass (Fig. 1). Each treatment and its respective controls were performed in triplicate over 10 days, using effluent and culture volumes as proposed by USEPA (1991).



**Fig. 1.** Graphical Representation of the Phycoremediation Assay.

### 2.3. Determination of Phosphorus and Nitrogen in Phytoremediation Treatments

Phosphorus and nitrogen concentrations were determined before and after exposure to the various treatments. Considering that microalgae can assimilate nitrogen in the form of nitrates (NO<sub>3</sub><sup>-</sup>), nitrites (NO<sub>2</sub><sup>-</sup>), and ammonium (NH<sub>4</sub><sup>+</sup>), and phosphorus as phosphate (PO<sub>4</sub><sup>3-</sup>), the percentage removal of these nutrients was recorded at the start (day 0), after 72 hours (day 3), and at the end (day 10). The following formulas were applied based on Lopez Espinoza (2022), with minor modifications for this experimental setup:

For Nitrates,

$$\left[ \left( \frac{\text{Nitrate (mg/L) Day 0} - \text{Nitrate (mg/L) Day 10}}{\text{Nitrate (mg/L) Day 0}} \right) \times 100 \right]$$

For Phosphates,

$$\left[ \left( \frac{\text{Phosphate (mg/L) Day 0} - \text{Phosphate (mg/L) Day 10}}{\text{Phosphate (mg/L) Day 0}} \right) \times 100 \right]$$

### 2.4. Cell Density Analysis

A 0.1 mL sample was taken from each replicate and placed on a microscope slide for observation under an optical microscope at 40x magnification. A Greek key pattern was used to traverse the slide, and individual cells within each filament were counted across the entire sample [20].

Cellular density was estimated by counting the total number of filaments per treatment (ft) in relation to the total number of filaments across all treatments (fT), calculating the relative frequency (Frt) as follows: [Frt = (ft / fT)] x 100. Similarly, the relative frequency of individual cells per treatment was estimated [21]. This data transformation into continuous variables

allowed fulfillment of statistical assumptions [21].

### 2.5. Morphological Analysis

Photographs of 10 fields per sample were taken under a 40x optical microscope using the preparations described in section 2.4. These images were analyzed using ImageJ software [22], calibrated with a millimeter scale to measure individual cell dimensions. These measurements were later used for statistical analysis to compare between groups.

### 2.6. Chlorophyll Determination

Chlorophyll values were estimated at wavelengths of 430 nm and 664 nm for chlorophyll a, 460 nm and 647 nm for chlorophyll b, and 442 nm and 630 nm for chlorophyll c, following Larkum (2003).

The following formulas from APHA (2012) were used:

Chlorophyll a:

$$[(11.85 \times 664 \text{ nm}) - (1.54 \times 647 \text{ nm}) - (0.08 \times 630 \text{ nm})]$$

Chlorophyll b:

$$[(21.03 \times 647 \text{ nm}) - (5.43 \times 664 \text{ nm}) - (2.66 \times 630 \text{ nm})]$$

Chlorophyll c:

$$[(24.52 \times 630 \text{ nm}) - (7.6 \times 647 \text{ nm}) - (1.67 \times 664 \text{ nm})]$$

### 2.7. Statistical Analyses

One-way ANOVA and Tukey's post hoc tests were used to compare nitrogen and phosphorus concentrations across treatments. The same statistical test was employed to evaluate algal growth among treatments. If data violated the assumptions of homogeneity of variances or normality, non-parametric Kruskal-Wallis tests with post hoc analysis were used. Additionally, principal component analysis (PCA) was performed to evaluate associations and responses of each analyzed variable to the treatments.

Finally, t-tests were used to compare phosphorus and nitrogen removal between the two types of treated effluents [21].

### 3. RESULTS

#### 3.1 Effluent Characterization

The following section presents the results of the physicochemical characterization of the effluent (Table 1), showing the initial values of phosphorus and nitrogen prior to the phycoremediation process.


**Table 1.** Data obtained from the determinations of physicochemical parameters of the urban effluent.

DETERMINATIONS	VALUES	USED METHOD
Turbidity - NTU	63	S.M. 4500-B
Suspended Solids (mg/L)	144.9	S.M. 2510-B
Conductivity ( $\mu$ S/cm)	750	S.M. 2540-B
Total Dissolved Solids – TDS (mg/L)	389	S.M. 2540-B
pH	7.01	S.M. 4500-B
Nitrate (mg/L)	28	S.M-4500-NO <sub>3</sub> <sup>-</sup> -E
Nitrite (mg/L)	0,16	S.M-4500-NO <sub>2</sub> <sup>-</sup> -B
Ammonium (mg/L)	1.5	S.M- 4500-NH <sub>4</sub> H
Sulfates (mg/L)	45	S.M-4500-SO <sub>4</sub> <sup>-2</sup> -E
Chlorides (mg/L)	24.9	S.M. 4500-Cl-B
Phosphorus (mg/L)	2.28	S.M- 4500-P-E
Chemical Oxygen Demand – COD (mg/L O <sub>2</sub> )	165.2	S.M. -5220-D
Biochemical Oxygen Demand – BOD (mg/L O <sub>2</sub> )	143.1	S.M. -5210-B

#### 3.2 Physicochemical Results of the Bioassays with Wastewater Effluents

##### 3.2.1 Phosphate Determination

The results showed a remediation efficiency of 100%, 97.17%, and 99.17% for the treatment with *M. varians* in effluent dilutions of 12.5%, 25%, and 50%, respectively. These results are shown in Fig. 2.




**Fig. 2.** Phosphorus removal by *Melosira varians* after 10 days. The treatments correspond to 12.5% (brown line), 25% (orange line), and 50% (green line) effluent dilution.

### 3.2.2 Nitrate Determination

The results indicated nitrate removal efficiencies of 29.41%, 41.18%, and

70.59% for the treatment with the microalga in effluent dilutions of 12.5%, 25%, and 50%, respectively.




**Fig. 3.** Nitrogen removal by *Melosira varians* after 10 days. The treatments correspond to 12.5% (brown line), 25% (orange line), and 50% (green line) effluent dilution.

### 3.3 Cell Density: Determination of Algal Growth

The following results show the growth of *M. varians* after 10 days of treatment. It is

worth noting that growth was assessed by measuring the number of cells and the number of filaments per treatment, as represented in Fig. 4. The ANOVA

statistical results showed no significant differences between treatments and the control group ( $p > 0.05$ ), despite a marked trend toward increased cell and filament numbers.




**Fig. 4.** Percentage of *M. varians* individuals after 10 days of exposure.

### 3.4 Cell Size: Determination of Cell Width and Length

#### 3.4.1 Cell Width Determination

The results of the cell width analysis evaluated in the controls and treatments are presented below. Statistical analyses

showed that 10 days after treatment, the width of the microalgae increased with increasing concentration of the effluent ( $p < 0.05$ ; Fig. 5).




**Fig. 5.** Cell width of *Melosira varians* measured in the control group and the effluent treatments at 12.5% (T1), 25% (T2), and 50% (T3), obtained on the tenth day of the bioassay (A).


### 3.4.2 Cell Length Determination

The results of the cell length analysis in the controls and treatments are presented below. Cell length values are shown in Fig. 6, while photographs illustrating these cellular changes are shown in Fig. 7.


Statistical analysis demonstrated that after 10 days of the bioassay, the length of the microalga increased in the 50% and 12.5% effluent treatments and decreased in the 25% effluent treatment ( $p < 0.05$ ; Fig. 6).



**Fig. 6.** Cell length of *Melosira varians* measured in the negative control group (CN) and the effluent treatments at 12.5% (T1), 25% (T2), and 50% (T3), obtained on the tenth day of microalga treatment (A).



**Fig. 7.** Optical microscope images (40X) of *Melosira varians* filaments corresponding to the control (A) and effluent treatments at 12.5% (B), 25% (C), and 50% (D) after 10 days of exposure.




**Fig. 8.** Scanning electron micrograph (SEM) of *Melosira varians*, showing the typical cylindrical shape of the genus.

### 3.5 Chlorophyll Determination

The results of the chlorophyll a, b, and c analyses in the controls and phycoremediation treatments are shown in Fig. 9.

After 10 days of phycoremediation, statistical analysis indicated that chlorophyll a levels showed no significant differences between the control and treatment groups ( $p > 0.05$ ; Fig. 9). However, a significant decrease in chlorophyll b was observed in the 25%

and 50% effluent treatments compared to the other groups ( $p < 0.05$ ; Fig. 9). Similarly, for chlorophyll c, a significant reduction was detected in the same effluent concentrations compared to the other treatments ( $p < 0.05$ ; Fig. 9). Finally, as observed in Fig. 9, chlorophyll c was the most abundant pigment across the various treatments, compared to chlorophyll a and b.



**Fig. 9.** Chlorophyll a (black bars), b (gray bars), and c (white bars) content in *Melosira varians* from the control (C) and the effluent treatments at 12.5% (T1), 25% (T2), and 50%

(T3) after 10 days of exposure. \* indicates statistically significant differences compared to the control and other treatments for the respective chlorophyll.

### 3.6. Multivariate Analysis: Principal Component Analysis

A positive association was observed between chlorophyll a and phosphate levels, as well as with chlorophylls b and c. Additionally, chlorophylls b and c showed a strong positive correlation with each other. Moreover, no correlation was found between chlorophyll content and biomass. This suggests a potential shift of the algae toward heterotrophic metabolism and nutrient uptake, where phosphorus is used for photosynthesis rather than biomass production. Conversely, biomass showed a positive correlation with cell width and length, highlighting the relevance of these measurements for estimating biomass in diatoms, as opposed to relying solely on chlorophyll content.

Furthermore, nitrate levels showed negative correlations with all analyzed biological parameters, suggesting a possible detrimental effect.

## 4. DISCUSSION AND CONCLUSION

Conventional wastewater treatment plants employ various processes to reduce the organic load in effluents. However, the effective removal of nitrogen and phosphorus remains a major challenge, often requiring additional treatment stages that are both energy-intensive and costly. This limitation underscores the need for alternative, sustainable methods capable of efficiently removing nutrients through environmentally friendly means. Such approaches are critical for the proper management of wastewater, as nutrient-rich discharges contribute not only to the eutrophication of surface and groundwater bodies but also to the

emission of greenhouse gases, posing a broader environmental threat [23].

The use of microalgae as an ecological alternative in tertiary wastewater treatment plants has attracted considerable interest in recent years. So far, the greatest advances in the application of this technology have been made with green algae (such as *Scenedesmus sp.* and *Chlorella sp.*) [24], [25]. However, we observe that the use of diatoms in phycoremediation of effluents offers significant advantages over green algae. Firstly, diatoms require less light than other algae; since their silica frustules are transparent, they can grow even on cloudy or rainy days and can also dominate under nutrient-limited conditions [26]. Additionally, it has been proven that under stress conditions, diatoms such as *Melosira varians* demonstrate the ability to assimilate significant amounts of nutrients, particularly nitrogen and phosphorus, which are essential for protein synthesis—accounting for 45–60% of microalgal dry weight [26]. When nutrients are abundant, these species commonly store energy in the form of carbohydrates or lipids, which serve as alternative energy sources during periods of darkness when photosynthesis is not feasible. This metabolic flexibility is critical for survival under fluctuating environmental conditions.

Diatoms are generally considered autotrophic organisms, relying primarily on light within the visible spectrum to conduct photosynthesis. However, recent studies have identified the presence of carbohydrate and amino acid transport mechanisms across several diatom species, suggesting a potential for facultative heterotrophy [27]. In benthic

species like *M. varians*, which are often located in microenvironments with limited light but abundant decomposing organic matter, the reliance on heterotrophic metabolism may be more common. This would explain their capacity to persist in dark conditions for extended periods [27], [28].

These characteristics—efficient nutrient assimilation, metabolic flexibility, and resistance to pollutants due to their silica frustules—make diatoms exceptionally well-suited for phycoremediation. *Melosira varians*, demonstrated notable efficacy in removing contaminants from domestic wastewater effluents, reinforcing its utility as a bio-remediator in aquatic systems.

With respect to pigment composition, the results of this study confirmed that *M. varians* possesses a higher proportion of chlorophyll c relative to chlorophylls a and b, consistent with diatom pigment profiles. Furthermore, the significant reduction in chlorophyll b and c in treatments with elevated nutrient loads may indicate a metabolic shift from autotrophy toward heterotrophy. In nutrient-stressed environments—particularly under nitrogen or silica limitation—diatoms rely on stored lipids as energy sources and suppress photosynthetic activity [29], [30]. This also accounts for the observed increase in cell size, which may result from the accumulation of lipid droplets within the cytoplasm.

The high nitrogen removal efficiency observed in the 50% effluent treatment (~70%) likely led to early nutrient depletion, placing the microalgae under physiological stress. This may have triggered a shift in metabolic strategy, consistent with a mixotrophic response. Our findings support the hypothesis that *M. varians* is capable of adjusting its nutritional mode depending on

environmental conditions. To our knowledge, this is the first report suggesting potential mixotrophy in *M. varians*.

Exposure of the algae to different concentrations of effluent showed different growth, responding favorably to higher concentrations of effluent [31].

Moreover, the decline in cell density observed under nutrient-limited conditions may be attributed to the initiation of a resting stage, possibly auxospore formation. This is a common strategy among diatoms to withstand unfavorable environments and may also contribute to increased cell volume [29], [30].

## 10. REFERENCES

1. Dana, Muñoz Carabajal, Daruich, Griselda Jorgelina, and Pedranzani, Hilda Elizabeth. 2024. "Utilizing Tetrademus Obliquus for Phycoremediation Enhanced Nitrogen and Phosphorus Removal in Urban Wastewater Treatment". *Journal of Global Ecology and Environment* 20 (4):169-79. <https://doi.org/10.56557/jogee/2024/v20i48955>.
2. Round, F. E., Crawford, R. M., & Mann, D. G. (1990). *The diatoms. Biology and morphology of the genera*. Cambridge University Press.
3. Smetacek, V. (1999). Diatoms and the ocean carbon cycle. *Protist*, 150(1), 25–32. [https://doi.org/10.1016/S1434-4610\(99\)70006-4](https://doi.org/10.1016/S1434-4610(99)70006-4)
4. Mann, D. G., & Vanormelingen, P. (2013). An inordinate fondness? The number, distributions, and origins of diatom species. *The Journal of Eukaryotic Microbiology*, 60, 414–420.
5. Khan, M. J., Raj, A., AHIRWAR, A., SIROTIYA, V., MOURYA, M., MISHRA, S., & VINAYAK, V. (2021). Diatom microalgae as smart

- nanocontainers for biosensing wastewater pollutants: Recent trends and innovations. *Bioengineered*, 12(2), 9531–9549.
6. Stevenson, F. (1996). Nitrogen-organic forms. In *Methods of Soil Analysis Part 3—Chemical Methods* (pp. 1185–1200).
  7. Wehr, J., & Sheath, R. G. (2003). *Freshwater algae of North America: Ecology and classification*. Academic Press.
  8. Azim, M. E., Verdegem, M. C. J., van Dam, A. A., & Beveridge, M. C. M. (2005). *Periphyton: ecology, exploitation and management*. CABI.
  9. Round, F. E. (1991). Diatoms in river water-monitoring studies. *Journal of Applied Phycology*, 3(2), 129–145.
  10. Biggs, B. J. F., & Kilroy, C. (2000). *Stream periphyton monitoring manual*. New Zealand Institute Water and Atmosphere for Ministry for the Environment.
  11. Potapova, M., & Charles, D. F. (2003). Distribution of benthic diatoms in U.S. rivers in relation to conductivity and ionic composition. *Freshwater Biology*, 48(8), 1311–1328.
  12. Marella, M. T., Bhaskar, M. V., & Tiwari, A. (2016). Phycoremediation of eutrophic lakes using diatom algae. En *Lake sciences and climate change* (pp. 103–115).
  13. Zhang, X. F., & Mei, X. Y. (2015). Effects of benthic algae on release of soluble reactive phosphorus from sediments: A radioisotope tracing study. *Water Science and Engineering*, 8(2), 127–131.
  14. Moreno, L. E. (2022). Ajuste local de índices que emplean diatomeas para determinar la calidad del ambiente fluvial serrano (Trabajo final correspondiente a la Carrera de Especialización en Calidad Ecológica y Restauración de Sistemas Fluviales, 44 págs.). Universidad Nacional de Luján.
  15. Lu, X., Liu, Y., & Fan, Y. (2020). Diatom Taxonomic Composition as a Biological Indicator of the Ecological Health and Status of a River Basin under Agricultural Influence. *Water*, 12(7), 2067. <https://doi.org/10.3390/w12072067>
  16. Gil, M. A., Daruich, G. J., Ulacco, J. H., & Moreno, L. E. (2020). Estudios sobre la calidad de los ambientes acuáticos de la región de Cuyo con aplicación de indicadores bióticos: Antecedentes y perspectivas. En E. Domínguez, A. Giorgi, & N. Gómez (Eds.), *La bioindicación en el monitoreo y evaluación de los sistemas fluviales de la Argentina: Bases para la integridad ecológica* (pp. 132–146). Eudeba.
  17. Guillard, R. R. (1975, October). Culture of phytoplankton for feeding marine invertebrates. In *Culture of marine invertebrate animals: proceedings—1st conference on culture of marine invertebrate animals greenport* (pp. 29-60). Boston, MA: Springer US.
  18. American Public Health Association (APHA). (2023). *Standard methods for the examination of water and wastewater* (24.<sup>a</sup> ed.). En W. C. Lipps, E. B. Braun-Howland, & T. E. Baxter (Eds.), APHA Press.
  19. Lucero, A. S. (2015). *Cultivo de consorcio microalgal autóctono y biorremediación de efluentes urbanos* [Tesis doctoral, Universidad Nacional de Cuyo].
  20. Villafañe, V. E., & Reid, F. M. H. (1995). *Métodos de microscopía para la cuantificación del fitoplancton. Manual de métodos ficológicos*. Universidad de Concepción, Concepción, 169-185.
  21. Zar, J. H. (2010). *Biostatistical analysis*. Pearson Education.
  22. Sunoj, S., Hammed, A., Igathinathane, C., Eshkabilov, S., & Simsek, H. (2021). Identification, quantification, and growth profiling of eight different microalgae species using image analysis. *Algal Research*, 60, 102487.

23. Garduño-Pineda, L., Linares-Hernández, I., Martínez-Miranda, V., Teutli-Sequeira, E. A., Santa Cruz, J. M., & García-Sánchez, J. J. (2023). Sustainable removal of nutrients (n and p) in a wastewater treatment plant, with eggshell (biocalcium). *Heliyon*, 9(11).
24. Pérez-Silva, K. R., Vega-Bolaños, A. M., Hernández-Rodríguez, L. C., Parra-Ospina, D. A., & Ballén-Segura, M. A. (2016). Uso de *Scenedesmus* para la remoción de metales pesados y nutrientes de aguas residuales de la industria textil. *Ingeniería Solidaria*, 12(20), 95–105.
25. Kaloudas, D., Pavlova, N., & Penchovsky, R. (2021). Phycoremediation of wastewater by microalgae: A review. *Environmental Chemistry Letters*, 19, 2905–2920.
26. Kiran, M. T., Bhaskar, M. V., & Tiwari, A. (2016). Phycoremediation of eutrophic lakes using diatom algae. En M. P. Sharma (Ed.), *Advances in algal research* (pp. 103–122). InTech. <https://doi.org/10.5772/64111>
27. Vilchis, M. C. L., Fuerte, F. O. L., & Rojas, C. A. P. (2020). Algas de Cristal; diatomeas.
28. Salati, S., D'Imporzano, G., Menin, B., Veronesi, D., Scaglia, B., Abbruscato, P., Mariani, P., & Adani, F. (2016). Mixotrophic cultivation of *Chlorella* for local protein production using agro-food by-products. *Bioresource Technology*, 203, 82–89.
29. Lee, R. (2018). *Phycology* (5.<sup>a</sup> ed.). Cambridge University Press. <https://doi.org/10.1017/9781316407219>
30. Domozych, D. S., & LoRicco, J. G. (2023). The extracellular matrix of green algae. *Plant Physiology*, 1–18. <https://doi.org/10.1093/plphys/kiad384>
31. Cervantes-Urieta, V. A., Pérez-Castro, D., Galeana-Parra, M. A., Ramírez-Fuentes, E., & Trujillo-Tapia, M. N. (2020). Cultivo y composición bioquímica de diatomeas marinas (Bacillariophyta) de la Bahía de Santa Lucía, Acapulco, México. *Gayana. Botánica*, 77(1), 11-22.
32. Acevedo, S., Peñuela, G., & Pino, N. (2017). Producción de biomasa de *Scenedesmus* sp. y eliminación de nitrógeno y fósforo en aguas residuales domésticas. *Ingeniería y Competitividad*, 19(1), 185–193.
33. Aydın, G., & Büyükişik, B. (2014). Efectos de los pulsos de nutrientes específicos de especies: *Thalassiosira allenii* (Takano). *Journal of Tekirdag Agricultural Faculty*, 11(3), 82–90.
34. Berner, F., Heimann, K., & Sheehan, M. (2015). Biofilms de microalgas para la producción de biomasa. *Journal of Applied Phycology*, 27, 1793–1804.
35. Castillo-Morales, G. (2004). *Ensayos toxicológicos y métodos de evaluación de calidad de aguas: estandarización, intercalibración, resultados y aplicaciones*. IMTA.
36. Chapman, R. (2013). Algae: The world's most important "plants"—An introduction. *Mitigation and Adaptation Strategies for Global Change*, 18, 5–12.
37. Daruich, G. J., Algozino, C., Almeida, C. A., Gallardo, M., & Pérez-Iglesias, J. M. (2023). Diatomeas bénticas de un río urbano de San Luis, Argentina. *XXXIX Jornadas Argentinas de Botánica, XIV Simposio Argentino de Ficología*.
38. Daruich, G. J., Gil, M. A., & Moreno, L. E. (2020). Estructura de la taxocenosis de diatomeas y su relación con variables ambientales en tres cuencas de la provincia de San Luis (Argentina). *Biología Acuática*.
39. Daruich, J. (2007). *Composición y distribución de las Bacillariophyceae de la cuenca del río Nogolí, San Luis, Argentina*

- [Tesis doctoral, Universidad Nacional de San Luis].
40. Daruich, J., De Giuseppe, B., Zapico, G., & Ulacco, J. (2021). Distribución estacional de las diatomeas de un río serrano: Su relación con las variables ambientales. *Cymbella*, 6(3).
  41. Daruich, J., Gonzalez, P., Almeida, C., & Pérez-Iglesias, J. M. (2020). Uso de la alga verde *Oocystis* sp. como modelo biológico para evaluaciones toxicológicas de BDE-209. *Biocell*, 1, 1–6.
  42. Daruich, J., Tripole, S., Gil, A., & Vallania, A. (2013). Comunidades de algas y cianobacterias en dos ríos de la provincia de San Luis (Argentina) sometidos a influencia antropogénica. *Acta Limnologica Brasiliensia*, 25(1), 79–90.
  43. De Giuseppe, B., & Daruich, J. (2019). Estudios preliminares de la ficoflora del suelo en Salinas del Bebedero, San Luis, Argentina. *Biocell*, 43(4), 299–303.
  44. Donlan, R., & Costerton, J. (2002). Biofilms: Survival mechanisms of clinically relevant microorganisms. *Clinical Microbiology Reviews*, 15(2), 167–193.
  45. García-Pichel, F., & Belnap, J. (2021). Cyanobacteria and algae. En *Principles and applications of soil microbiology* (pp. 171–189).
  46. Jimenez Veuthey, M., Zapata, L. M., & Flores, M. L. (2020). Tratamiento de un efluente agroindustrial empleando la microalga de agua dulce *Scenedesmus obliquus* autóctona del Embalse Salto Grande (Entre Ríos, Argentina). *Revista Latinoamericana de Biotecnología Ambiental y Algal*, 11(1), 34–52.
  47. Juárez, A. B. (2017). Exposición a contaminantes de diferente naturaleza (herbicidas y antibióticos) sobre marcadores morfológicos, ultraestructurales y bioquímicos en poblaciones y comunidades de microalgas. En *Algas, helechos y plantas vasculares acuáticas en estudios de contaminación ambiental y ecotoxicología*. Universidad Nacional de Mar del Plata.
  48. Larkum, A. W. D., Douglas, J. A., & Raven, J. A. (2003). *Photosynthesis in algae*. Springer Science & Business Media.
  49. López, F. O., & Altamirano, M. C. (2011). Diatomeas bentónicas de los oasis de Baja California Sur [Informe Final SNIB-CONABIO, Proyecto No. HJ032]. Universidad Autónoma de Baja California Sur.
  50. Marella, T. K., López-Pacheco, I. Y., Parra-Saldívar, R., Dixit, S., & Tiwari, A. (2020). Wealth from waste: Diatoms as tools for phycoremediation of wastewater and for obtaining value from the biomass. *Science of the Total Environment*, 724, 137960.
  51. Marella, T. K., Saxena, A., & Tiwari, A. (2020). Diatom mediated heavy metal remediation: A review. *Bioresource Technology*, 305, 123068. <https://doi.org/10.1016/j.biortech.2020.123068>
  52. Mehrabadi, A., et al. (2015). Tratamiento de aguas residuales de alta tasa de estanques de algas (WWT HRAP) para la producción de biocombustibles de bajo costo. *Biosource Technology*, 184, 202–214.
  53. Mengo, L. (2017). Composición taxonómica y riqueza de diatomeas en el río Suquía, Córdoba (Argentina). *Revista Facultad de Ciencias Exactas, Físicas y Naturales*, 4, 55–62.
  54. Mohsenpour, M., et al. (2021). Integrating micro-algae into wastewater treatment: A review. *Science of the Total Environment*, 752, 142168.
  55. Muñoz, D. (2024). *Remoción de nutrientes de efluentes urbanos por fitorremediación con la microalga local Scenedesmus sp.*
  56. Olgún, E. J. (2003). Phycoremediation: Key issues for cost-effective nutrient removal processes. *Biotechnology Advances*, 22(1–2), 81–91.

57. Park, J. B. K., Craggs, R. J., & Shilton, A. N. (2011). Wastewater treatment high-rate algal ponds for biofuel production. *Bioresource Technology*, *102*(1), 35–42.
58. Pérez, D., & Menone, M. (2017). *Algas, helechos y plantas vasculares acuáticas en estudios de contaminación ambiental y ecotoxicología* (1.<sup>a</sup> ed.). Universidad Nacional de Mar del Plata. ISBN 978-987-544-756-1.
- 59.
60. Pérez-Vázquez, A., et al. (2024). Integrative assessment of in situ combined bioremediation strategies applied to remediate soils spilled with sewage sludges. *Frontiers in Environmental Science*, *12*, 1370820.
61. Pirjo, H., & Gómez, I. (2017). *Algas en ecotoxicología: Hacia el desarrollo de microbioensayos*. Instituto de Ciencias Marinas y Limnológicas, Facultad de Ciencias, Universidad Austral de Chile, Valdivia.
62. Prajapati, S. K., Kaushik, P., Malik, A., & Vijay, V. K. (2013). Phycoremediation coupled production of algal biomass, harvesting, and anaerobic digestion: Possibilities and challenges. *Biotechnology Advances*, *31*(8), 1408–1425.
63. Ray, A. M., Daruich, G. J., Algozino, C., Cortez, A., Jorquera, A., Muñoz, D., Quaife, E., Almeida, C., Gonzalez, P., Da Silva, S. M., Funes, M., & Pérez Iglesias, J. M. (2023). Toxicological bioassays on the native microalgae *Dictyosphaerium sp.* as a possible phycoremediator against relevant pesticides. *SETAC Latin America 15th Biennial Meeting, 17–20 September 2023, Montevideo, Uruguay*.
64. Rivas, M., & Riquelme, C. (2012). Probiotic biofilms. En *Probiotics*. IntechOpen.
65. Saxena, A., Tiwari, A., Kaushik, R., Iqbal, H. M., & Parra-Saldívar, R. (2021). Diatoms recovery from wastewater: Overview from an ecological and economic perspective. *Journal of Water Process Engineering*, *39*, 101705.
66. Sisman-Aydin, G. (2022). Comparative study on phycoremediation performance of three native microalgae for primary-treated municipal wastewater. *Environmental Technology & Innovation*, *28*, 102932.
67. Tong, C. Y., & Derek, C. J. (2021). Biofilm formation of benthic diatoms on commercial polyvinylidene fluoride membrane. *Algal Research*, *55*, 102260.
68. U.S. Environmental Protection Agency. (1991). *Methods for measuring the acute toxicity of effluents and receiving waters to freshwater and marine organisms* (p. 197). Environmental Monitoring Systems Laboratory, Office of Research and Development.
69. Viau, V. E., Marciano, A., Iriel, A., & López Greco, L. S. (2016). Assessment of a biofilm-based culture system within zero water exchange on water quality and on survival and growth of the freshwater shrimp *Neocaridina heteropoda heteropoda*. *Aquaculture Research*, *47*(8), 2528–2542.
70. Viau, V. E., Ostera, J. M., Tolvía, A., Ballester, E. L., Abreu, P. C., & Rodríguez, E. M. (2012). Contribution of biofilm to water quality, survival, and growth of juveniles of the freshwater crayfish *Cherax quadricarinatus* (Decapoda, Parastacidae). *Aquaculture*, *324*, 70–78.