

Phenotypic Resistance pattern of Bacterial Isolates from Gold Mine tailings in Ile Ife, Osun State, Nigeria

Abstract

Artisanal gold mining and the resultant environmental degradation and health hazards have been a cause for public health concern. This, specifically as heavy metal contaminated environments have been pointed out as major players in the dissemination of resistant pathogens of public health concern. This study aimed to determine the phenotypic antibiotic resistance profiles of bacteria isolated from gold mine tailings in a community in Osun State, Nigeria. Bacteria were isolated from excavation soil samples using standard microbiological methods, with general-purpose and selective media. Antibiotic susceptibility testing was performed by Kirby-Bauer disc diffusion method. Isolates recovered were identified and their zone of growth inhibition was determined. Of the 32 isolates identified, 100% were resistant to ceftriazone, and also 100% resistance respectively to oxacillin and amoxicillin-clavulanic acid by all the isolates tested, while 96.67% were resistant to ceftazidime. The isolates exhibited 12 phenotypic resistance pattern of which 100% were resistant to at least three types of antibiotics and over 45% MDR-multidrug resistance. These findings indicate a high frequency occurrence of antibiotic resistance in the goldmine tailings including multidrug resistant bacteria in the environment. Hence, this calls for the concerned stakeholders to take necessary measures in safeguarding the catchment community against this public health menace.

Key Words: Gold mining, Antibiotic, MDR-multidrug resistance, Environment

1. INTRODUCTION

Antibiotics resistance has continued to be a global public health concern that could lead to a post-antibiotic era, in which even minor infections would be life-threatening, as declared by World Health Organization [1]. Despite measures to counter this phenomenon, antibiotic

resistance in pathogens is still prevalent [2]. This implies the presence of extraneous factors, other than the antibiotics, that may be responsible for the evolution, selection and dissemination of antibiotic-resistant strains. In that regard, antimicrobial resistance (AMR) in environmental micro-organisms has been pointed out as a major culprit [3,4].

Heavy metals can co-select for antimicrobial resistance by fostering environments where resistance genes, whether within closely related taxa or across genera, undergo horizontal transfer [5,6]. Heavy metal(HM) contaminants have also been implicated in selection of microbial virulence factors and with the potential to alter the antibiotic resistome of contaminated water bodies [7,8]. Exposure to HM predisposes bacteria to antibiotic resistance, as both phenomena create selection pressure on the microbes [6, 9]. It has been established that in heavy metal contaminant ecosystem, bacteria can exhibit selection for heavy metal resistance and antibiotic resistance simultaneously [5]. This phenomenon could be by either co-selection when there are two or more genetically related resistance genes or cross-selection due to presence of single genetic element which provides tolerance to more than one antimicrobial agents [6,10].

Extrachromosomal resistance is commonly implicated in AMR in environmental microbes. These resistance determinants are transferrable to clinical pathogenic bacteria, as have been previously documented [2]. Therefore, researchers are focusing on the effects of mining on the environment, among which is the impact and relationship between HM contamination and antibiotics resistance [2, 6, 11].

Although gold mining provides huge socio-economic benefits to nations, the long-term deleterious impact on the environment and the public health cannot be overlooked [12,13]. Large quantities of waste are produced during gold mining activities, during which over 99% of the extracted ore is released as waste to the environment [14].The type of mining process applied in gold extraction and the quantity of waste generated is determined by the location of the gold deposit [15]. Mine tailings accumulate either in the form of large hills or water ponds, which could constitute severe environmental problems [16].

In Nigeria, gold mining is predominantly artisanal, carried out through open-pit methods that generate substantial environmental waste and heavy metal contamination, with adverse health implications [17,18,19]. Previous studies have examined the environmental impacts of these

activities, including contamination of water, soil and vegetation [20,21], yet data on the phenotypic antibiotic resistance profiles of soil bacteria in mining areas remain scarce. This study investigates the phenotypic antibiotic resistance profiles of bacteria isolated from gold mine tailings in IloroAraromi village, Ile-Ife, Osun State, Nigeria, to provide insights into the potential public health risks posed by these contaminated environments.

2. MATERIALS AND METHODS

2.1 Study area

The study area is IloroAraromi village originally known as AraromiOkeodo village situated in Kere area of Ife South (Fig 1), Osun State, Nigeria and its geographical coordinates are 7° 9' 9" North, 4° 27' 14" East. The climate is tropical, farming combines coco and banana plantation by residents of the community.

2.2 Sample collection

Soil samples were aseptically collected from mine tailings with a sterile spatula into sterile screw capped bottles. Composite samples were obtained by collecting at different excavations sampling points from top to a depth of 0-10cm below the top soil surface from two separate points respectively [15]. The soil samples were transported immediately to the laboratory for microbiological analysis.




Figure 1: Maps highlighting the study area. (A) Map of Nigeria displaying the sample location base on region (B) Map of Osun State showing the sample location (C) Outlined map of the study area

2.3 Preparation of media

Eosin methylene blue agar (EMBA), *SalmonellaShigella* agar (SSA) and Nutrient agar (NA) were prepared according to the manufacturers' instruction and were sterilized by autoclaving at 121°C for 15mins. SSA was sterilized by bringing to boil on a laboratory hot plate. All sterile media were poured into petri dishes at about 45°C and allowed to solidify. EMBA plates were used to isolate *E. coli* and other Gram-negative enteric bacteria, while SSA plates were solely for isolation of *Shigella* and *Salmonella* species. All media except otherwise stated were purchased from Hi-Media laboratories (Mumbai, India).

2.4 Bacterial isolation

Serial dilution of each soil sample was carried out using normal saline as diluent in test tubes containing 9ml of sterile water each. A 1 g of the soil sample was dispensed in a diluent test tube and homogenized. Thereafter, 1ml suspension of the homogenate was transferred using a sterile pipette into another diluent test tube and swirled gently to make 10^{-1} dilution. This procedure was repeated for the successive dilutions up to 10^{-6} dilution. One milliliter from dilutions 10^{-4} to 10^{-6} was inoculated in various culture media by spread plate method. However, for the isolation of *Salmonella* and *Shigella* 1ml 10^{-1} and 10^{-2} dilutions were inoculated into petri dishes then the prepared SSA was introduced and the plates were gently swirled to ensure even distribution of the inoculum. All inoculation of dilutions was in duplicate plates. The inoculated and blank control plates were incubated in an inverted position for 24hrs at 37°C in an incubator. The colonies that emerged were counted and colony forming units per ml was calculated. Sub culturing of colonies was done and pure cultures obtained were stored at 4°C in NA slants for further study.

2.5 Bacterial identification and characterization

Gram staining and biochemical tests were performed to differentiate organisms based on the structure of their cell wall and metabolic characteristics respectively. The biochemical tests comprised catalase, coagulase, IMViC (Indole, Methyl red, Voges proskauer) tests.

2.6 Antibiotics Sensitivity Testing

The isolates from gold mine soil were tested for antimicrobial susceptibility by disc diffusion using the Kirby Bauer disk diffusion method on Muller Hilton agar (BIOTEC) was performed with slight modification as described by [22], and Clinical and Laboratory Standards Institute [23] guideline was used to interpret the result. The following antibiotics with their different concentrations were used for testing: Ceftazidime ($30\mu\text{g}$), Gentamicin ($10\mu\text{g}$), Oxacillin ($1\mu\text{g}$), Ciprofloxacin ($5\mu\text{g}$), Ceftriazone ($30\mu\text{g}$) and Amoxicilin-clavulanic acid ($30\mu\text{g}$). *E. coli* ATCC 25922 and *S. aureus* ATCC 25923 served as the control strains. The definition of multidrug resistance (MDR) was in accordance with Centre for Disease Control (CDC) for resistance of bacterial isolates to at least one antibiotic in three or more drug classes [24].

2.7 Statistical Analysis

Data generated from bacterial isolation and antibiotic sensitivity tests were analyzed by entering into Microsoft Excel to determine the frequency of occurrence and resistance profile of the bacterial isolates using descriptive statistics.

3. RESULT AND DISCUSSION

3.1 Total heterotrophic and coliform bacterial count

The total bacterial population of the gold mine excavation soil recorded during the study is presented in Table 1. Heterotrophic plate count from the soil samples showed a highest value of 1.2×10^6 CFU/g observed at site C2, followed by 9.6×10^5 in site C1, while the lowest was 3.3×10^5 recorded at site CP1. Furthermore, the highest coliform count of 8.4×10^2 CFU/g was also documented at site C2 while the least count of 1.1×10^2 was observed at site C1. Comparatively, a previous study on a mining site in north-west Nigeria recorded highest and lowest bacterial counts of 4.5×10^6 cfu/g and 3.3×10^4 CFU/g respectively [25]. A low microbial density is usually associated with presence of heavy metal [26] which could explain the low HTB count recorded in this study. This is because, although needed in trace amounts, at higher concentrations heavy metals become toxic to microbes [27]. Nevertheless, the presence of coliform in the mining sites is a cause for environmental health concern as vast amounts of the contaminated soil can be spread to the community via aeolian and water erosion [28].

S/N	Site Code	Total heterotrophic count CFU/g	Total coliform count CFU/g
1	C1	9.6×10^5	1.1×10^2
2	C2	1.2×10^6	8.4×10^2
3	CP1	3.3×10^5	7.1×10^2

excavation soil

3.2 Bacterial identification and frequency rate

Thirty-two bacteria were isolated from the goldmine soil of which 26 (81%) were Gram negative and 6 (19%) were Gram positive. A total of 10 genera were identified, and the highest frequency rate of 16% recorded was for *Klebsiella*, *Proteus* and *Citrobacter* species respectively. This is followed by *Providencia* and *Staphylococcus* species with 13 and 12% frequency rate respectively, while the least was *Edwardsiella* sp. with 3% frequency rate. The result is depicted in Fig. 2. This result implies that gold mine excavations in the studied area harbour a number of enteric bacterial genera constituting public health threat as opportunistic pathogens. This is probably due to ‘bush method’ of defecation and human waste management system, common in many rural and peri-urban communities [29]. Specifically, *Citrobacter* spp. occur in animal and human guts and in some environment including soil and sewage sludge water, but have also been implicated in 3-6% of all *Enterobacteriaceae* isolates responsible for hospital acquired infections [30]. In addition, *Proteus*, *Klebsiella* and *Providencia* species although occurring in natural environment, are capable of colonizing skin and mucosal tissues. These have been implicated as agents of healthcare-associated infections (HAIs) such as urinary tract infection and burn wounds as well as gastroenteritis, including travelers' diarrhea [31]. A report on AMR status in WHO African regions stated that *Klebsiella* spp. were the most common bacterial infectious agents in Africa [32]. Noteworthy also is the incidence (6% respectively) of *E. coli* and *Salmonella* spp. recorded in this study, which are common agents of enteric fever and gastrointestinal disorders. For instance, illegal miners were reported to be the predominant group in an enteric fever

outbreak in South Africa probably due to consumption of contaminated ground water while working in a gold mine shaft [33].

Previous studies have reported several genera and bacterial phyla in mining sites; Proteobacteria, Actinobacteria, and Firmicutes were reported as the predominant phyla in abandoned gold mine tailings in Krugersdorp, South Africa [15]. Furthermore, other studies on HM contaminated environments in Nigeria and some other countries have reported different bacterial genera including those carrying HM-resistant genes such as *Bacillus* and *Gemmata*[34], *Pseudomonas* and *Bacillus*[35], *Priestiaaryabhatai* and *Enterobactercloacae*[36].

However, it should be noted that frequency and diversity of microbial population in HM contaminated sites are a function of the HM concentration and the microbes that have been able to adapt to the prevailing conditions in such environment [5,27, 37].




Fig. 2. Frequency rate of bacterial genera isolated from gold mine excavation soil

3.3 Resistance profile of bacterial isolates

All bacterial isolates in this study demonstrated complete resistance (100%) to ceftriaxone, oxacillin, and amoxicillin-clavulanic acid, with 96.67% also resistant to ceftazidime (Fig. 3). On the other hand, gentamicin and ciprofloxacin retained susceptibility in 84.38% and 65.63% of the isolates, respectively. The high resistance to first-line antibiotics such as ceftriaxone and amoxicillin-clavulanic acid, both widely used in primary and emergency care globally, is particularly concerning, indicating limited treatment options for infections by these isolates [38]. Oxacillin, is a β -lactam antibiotic member also referred to as penicillinase resistant penicillin

being resistant to β -lactamase hydrolytic enzyme. The antibiotic has a limited spectrum of activity mostly on Gram-positive bacteria with resistance being common particularly among *Enterobacteriaceae* [39]. Conversely, ceftazidime and ceftriazone are third-generation cephalosporin antibiotics used in the management of most Gram-negative bacterial infections, especially those resistant to first- and second-generation cephalosporin or other β -lactam antimicrobials [40]. Moreover, amoxicillin-clavulanic acid (β -lactam/ β -lactamase inhibitor combinations) is the most frequently used antimicrobial combination by primary care givers and in emergency conditions globally [38]. Amoxicillin is a broad spectrum β -lactam penicillin derivative with similar activity against Gram- positive and negative bacteria by inhibiting cell wall biosynthesis. The clavulanic acid is a β -lactamase inhibitor that prevents microbial degradation of the β -lactam, hence the combination offers extended antibiotic coverage and reduce resistance [38]. Nevertheless, the high resistance observed in this study suggests that reliance on these antibiotics for managing infections linked to this environment may be ineffective, underlining the need for continuous antimicrobial resistance surveillance. Major mechanisms of β -lactam resistance include resistance by β -lactamase enzyme hydrolysis of the antibiotic, resistance by active efflux of the antibiotic and resistance by penicillin-binding protein modification and increased peptidoglycan synthesis [39].




Fig. 3: Antibiotics Sensitivity Profile of Bacterial Isolates from Gold Mine Excavation Soil

UNDER PEER REVIEW

Table 2: Phenotypic resistance pattern of bacteria isolated from gold mine excavation soil

No. of Antibiotics	Resistance Phenotype	No. of isolates (%) with resistance	Bacteria	
3	OX, CRO, CAZ	2(6.3%)	Kleb(BB1) and Staph(BB2)	
	AMC, CRO, CAZ	2(6.3%)	Sal(BB3) and Prov(B2b)	
	OX, AMC, CRO	2(6.3%)	Prov(B8) and Prot(B10b)	
4	OX, AMC, CRO, CAZ	1(3.1%)	Staph(BP5)	
	AMC, CRO, CAZ, CIP	2(6.3%)	Prot(BB4) and Prov(A1)	
	OX, CRO, GN, CAZ	1(3.1%)	Edws(BP3)	
	OX, AMC, CRO, CIP	1(3.1%)	Prot (A3)	
	OX, CRO, CAZ, CIP	2(6.3%)	Sal(BB5) and Staph(AP3)	
				Kleb(AB4), Entero(B7), <i>E. coli</i> (AP4) and (BP2), Prot(B6), Entero(A6), Prov(A7), Kleb(A9), Citro(B1a) and (B1b)
	OX, AMC, CRO, CAZ	10(31.3%)		
5	OX, AMC, CRO, GN, CAZ	3(9.4%)	Bac(A10), Kleb(AB5) and Citro(A5)	
	OX, AMC, CRO, CAZ, CIP	5(15.6%)	Staph(BP4), Citro(AB3), Prot(A2), Kleb(A8) and Bac(B5)	
	AMC, CRO, GN, CAZ, CIP	1(3.1%)	Citro(AP5)	

Key: Kleb-*Klebsiella*; Staph-*Staphylococcus*; Sal-*Salmonella*; Prov-*Providencia*; Prot-*Proteus*; Edws-*Edwardsiella*; Entero-*Enterobacter*; Citro-*Citrobacter*; Bac-*Bacillus*.

3.4 Multidrug resistance (MDR)

The isolates recovered from the gold mine excavation soils exhibited twelve (12) phenotypic resistance pattern (Table 2) of which 100% were resistant to at least three types of antibiotics while 31.3% were resistant to four (4) types of antibiotics comprising oxacillin, amoxicillin-clavulanic acid, ceftriazone and ceftazidime. A 46.88% multidrug resistance (MDR) comprising eight bacterial genera (Table 3) was also recorded. Furthermore, resistance to the highest number of antibiotics tested in this study was typical for *Klebsiella* spp. followed by *Citrobacter* spp. Likewise, the two *E. coli* isolates in this study were resistant to all the β -lactam class and the third-generation cephalosporins (Table 3), corroborating an earlier report by WHO which documented significant *E. coli* resistance to third generation cephalosporins and fluoroquinolones in communities and hospitals in African region [1]. Previous studies on bacterial isolates from heavy-metal contaminated freshwater farms, seawater farms, and their

markets, and dumpsite respectively reported 40 and 50% MDR [41,42]. This is somewhat similar to that of [8] in which bacteria phyla Proteobacteria, Firmicutes and Bacteroidetes were 87 and 50% resistant to at least one antibiotic and more than three antibiotic classes respectively. On the other hand, *Staphylococcus* spp. was reported as having high tolerance to several antibiotics tested for heterotrophic bacteria isolated from mine tailings in Poland [43].

Table 3: Multidrug resistant bacterial isolates from goldmine tailings

No. of Antibiotics	Resistance Phenotype	No. of isolates (%)	Bacteria
MDR	AMC, CRO, CAZ, CIP	2	Prot(BB4) and Prov(A1)
	OX, CRO, GN, CAZ	1	Edws(BP3)
	OX, AMC, CRO, CIP	1	Prot (A3)
	OX, CRO, CAZ, CIP	2	Sal(BB5) and Staph(AP3)
	OX, AMC, CRO, GN, CAZ	3	Bac(A10), Kleb(AB5) and Citro(A5)
	OX, AMC, CRO, CAZ, CIP	5	Staph(BP4), Citro(AB3), Prot(A2), Kleb(A8) and Bac(B5)
	AMC, CRO, GN, CAZ, CIP.	1	Citro(AP5).
Total number of isolates (%)		15(46.88%)	

Key: Kleb-*Klebsiella*; Staph-*Staphylococcus*; Sal-*Salmonella*; Prov-*Providencia*; Prot-*Proteus*; Edws-*Edwardsiella*; Citro-*Citrobacter*; Bac-*Bacillus*; MDR-Multidrug resistance.

Empirical administration of antibiotics is a common practice in most African countries, in which antibiotics are prescribed for patients without antibiotic sensitivity testing and in non-bacterial infections including cases of COVID-19 [44]. In Nigeria as in many other developing countries, antibiotics are unregulated, cheap and easily accessible over the counter without prescription [44]. This is despite the WHO established guidance on the appropriate use of antibiotics for common infections [45]. It is noteworthy that the result of our study indicates high resistance to the Access and Watch antibiotic groups comprising those recommended by WHO for the empiric treatment of most common infections, are less costly and have lower potential for the selection of antimicrobial resistance, and the ones that the use should be carefully monitored to avoid overuse due to a higher potential for the selection of antimicrobial resistance respectively [45]. Therefore,

the resistance pattern in this study implies the types of antibiotics usage in the community, since non-metabolized antibiotics can be discharged into the environment [27]. The multidrug resistant bacterial genera recovered from the mining sites could serve as potential transmission channel to the nearby community.

4. CONCLUSION

This study highlights the significant presence of antibiotic-resistant bacteria including multidrug resistant bacteria in gold mining soils from IloroAraromi village, Ile-Ife, Osun State, Nigeria. The findings revealed a high prevalence of resistance to commonly used antibiotics suggesting that environmental exposure in heavy metal-contaminated soils may contribute to the selection and spread of resistant bacteria. These findings call for urgent intervention to the environmental and public health impacts of artisanal mining practices, as well as the need for targeted surveillance and mitigation strategies to control the spread of antibiotic resistance in such environments. Broader regional sampling and molecular techniques are recommended to better understand the links between heavy metal contamination and antimicrobial resistance development in mining areas.

References

1. WHO (2014) *WHO's first global report on antibiotic resistance reveals serious, worldwide threat to public health*. Available at: <https://www.who.int/news/item/30-04-2014-who-s-first-global-report-on-antibiotic-resistance-reveals-serious-worldwide-threat-to-public-health> (August, 2022)
2. Vats P, Kaur UJ, Rishi P. Heavy metal-induced selection and proliferation of antibiotic resistance: A review. *J. Appl Microbiol.* 2022;132, 4058-4076
3. Fletcher S. Understanding the contribution of environmental factors in the spread of antimicrobial resistance. *Environ. Health Prev Med.* 2015;20(4):243-52. <https://doi:10.1007/s12199-015-0468-0>.

4. Ahmad SI, Malak HA, Abulreesh HH. Environmental antimicrobial resistance and its drivers: a potential threat to public health. *J. Global Antimicrob. Res.* 2021;27:101-111. <https://doi.org/10.1016/j.jgar.2021.08.001>
5. Chen J, Junjian L, Zhang H, Shi W, Liu Y. Bacterial Heavy-Metal and Antibiotic Resistance Genes in a Copper Tailing Dam Area in Northern China. *Frontiers in microbiology.* 2019;10:1916.
6. Sun F, Xu Z, Fan L. Response of heavy metal and antibiotic resistance genes and related microorganisms to different heavy metals in activated sludge. *J. Environ. Manag.* 2021;300:113754. <https://doi.org/10.1016/j.jenvman.2021.113754>
7. Lear L, Hesse E, Newsome L, Gaze W, Buckling A, Vos M. The effect of metal remediation on the virulence and antimicrobial resistance of the opportunistic pathogen *Pseudomonasaeruginosa*. *BioRxiv.* 2022; <https://doi.org/10.1101/2022.09.20.508257>
8. Bosch J, Bezuidenhout C, Coertze R, Lesego M-T. Metal- and antibiotic-resistant heterotrophic plate count bacteria from a gold mine impacted river: the Mooi River system, South Africa. *Environ. Sci. Pollution Res.* 2023;30:31605–31619. <https://doi.org/10.1007/s11356-022-24015-3>
9. Bezuidenhout CC, Mienie CC, De Klerk T, Molale L. Geospatial analysis of microbial community structure and antimicrobial resistance analysis in the management of natural streams and selected wetlands. Report to the Water Research Commission, WRC Report 2017; No k5/2347//3, WRC Report No k5/2347//3
10. Perelomov L, Sizova O, Gertsen M, Perelomova I, Arlyapov V, Atroshchenko Y. Antibiotic Resistance in Metal-Tolerant Microorganisms from Treatment Facilities. *Antibiotics.* 2023;12(12), 1678.
11. Hu HW, Wang JT, Li J, Li JJ, Ma YB, Chen D, He JZ. Field-based evidence for copper contamination induced changes of antibiotic resistance in agricultural soils. *Environ. Microbiol.* 2016;18(11), 3896-3909.
12. Environmental Law Institute, ELI. Artisanal and Small-Scale Gold Mining in Nigeria: Recommendations to Address Mercury and Lead Exposure. Environmental Law Institute®, Washington, D.C. ELI Project No. 12100. 2013; <https://www.eli.org/sites/default/files/eli-pubs/nigeria-asgm-assessment-final-report.pdf>

(Accessed April 2024).

13. Ondayo MA, Watts MJ, Mitchell CJ, King DCP, Osano O. Review: Artisanal Gold Mining in Africa—Environmental Pollution and Human Health Implications. *Exposure and Health*.2023;<https://doi.org/10.1007/s12403-023-00611-7>(Accessed April 2024).
14. Se A, Bo O, Fa O, Akinnifesi O. Assessment of heavy metal pollution in a gold mining site in Southwestern Nigeria. *Biomed. J. Sci. Tech. Res.*2018;12.<https://doi.org/10.26717/BJSTR.2019.12.002276>
15. Fashola MO, Ngole-Jeme VM, Babalola OO.Physicochemical properties, heavy metals, and metal-tolerant bacteria profiles of abandoned gold mine tailings in Krugersdorp, South Africa.*Can. J. Soil Sci.*2020;100: 217–233. <https://dx.doi.org/10.1139/cjss-2018-0161>.
16. Jiang X, Liu W, Xu H, Cui X, Li J, Jurong. Characterizations of heavy metal contamination, microbial community, and resistance genes in a tailing of the largest coppermine in China. *Environ. Pollution*.2021;280:116947. <https://doi.org/10.1016/j.envpol.2021.116947>
17. Ako TA, Onoduku US, Oke SA, Adamu IA, Ali SE, Mamodu A, Ibrahim AT.Environmental Impact of Artisanal Gold Mining in Luku, Minna, Niger State, North Central Nigeria. *J Geosci Geomatics*.2014;2(1):28-37.
18. Peter BA, Zakaria OM, Sabiki SA. Environmental and Health Impact of Small-Scale Gold Mining Activities in Ibanda District of Uganda. *J. Appl. Sci. Environ. Manage.* 2022;26(5):785-788.
19. Oluwafemi O, Akinwumiju A, Oladepo O, Oladepo D. Assessment of Gold Mining Impact on the Vegetation of Itagunmodi-IgunDrainage Basin, Southwestern Nigeria from 1972-2011: A Remote Sensing. *Proceedings of the International Cartographic Association*, 4, 2021; 30th International Cartographic Conference (ICC2021), Florence, Italy.<https://doi.org/10.5194/ica-proc-4-78-2021> (Accessed April 2024).
20. Ayantobe OO, JA, Awomeso GO, Oluwasanya BS, Bada AM, Taiwo. Gold mining in Igun-Ijesha, Southwest Nigeria: impacts and implications for water quality.*Amer. J. Environ. Sci.*2014;10(3):89-300.<https://doi:10.3844/ajessp.2014.289.300>.

21. Adekiya AO, Oloruntoba A, Ewulo BS, Ogunbode T, Sangoyomi T. Heavy Metal Pollution in Soils near Ilesha Gold Mining Area, Nigeria. *The Open Agric.* 2024;18:e18743315290216. <https://doi.org/10.2174/0118743315290216240123111021>
22. Fatokun EN, Fakorede CN, Atobatele KZ. Antibiotic Susceptibility Profile of Bacterial Isolates from Commercial Poultry Farms in Ile-Ife, Nigeria. *Chem. Biomol. Eng.* 2021;6(3):59-67. <https://doi.org/10.11648/j.cbe.20210603.13>
23. CLSI. Performance Standards for Antimicrobial Susceptibility Testing. 2017;27th Edition. Wayne, USA.
24. CDC. (<https://www.cdc.gov/narms/resources/glossary.html>). (Accessed March 2022)
25. Ibrahim UB, Kawo AH, Yusuf I, Yahaya S. Physicochemical and molecular characterization of heavy metal-tolerant bacteria isolated from soil of mining sites in Nigeria. *J. Genetic Eng. Biotech.* 2021;19:152 <https://doi.org/10.1186/s43141-021-00251-x>
26. Ji H, Zhang Y, Bararunyeretse P, Li H. Characterization of microbial communities of soils from gold mine tailings and identification of mercury-resistant strain. *Ecotox. Environ. Safety.* 2018;165:182-193 <https://doi.org/10.1016/j.ecoenv.2018.09.011>.
27. Fashola MO, Anagun OS, Babalola OO. Heavy metal pollution: toxic effects on bacterial cells. Pre-print. 2023; <https://doi.org/10.22541/au.168690270.01555938/v1>
28. Callender K. The revegetation (phytostabilization) of gold mine residues at Val d'Or, QC using *Frankia*-inoculated alders. 2014; McGill University (Canada).
29. Twinomucunguzi FRB, Nyenje PM, Kulabako RN, Semiyaga S, Foppen JW, Kansime F. Reducing Groundwater Contamination from On-Site Sanitation in Peri-Urban Sub-Saharan Africa: Reviewing Transition Management Attributes towards Implementation of Water Safety Plans. *Sustainability.* 2020;12:4210. <https://doi.org/10.3390/su12104210>
30. Jabeen I, Islam S, Hassan AKMI, Tasnim Z and Shuvo SR. A brief insight into *Citrobacter* species- a growing threat to public health. *Front. Antibiot.* 2023;2:1276982. <https://doi.org/10.3389/frabi.2023.1276982>.
31. dos Santos GS, Solidônio EG, Costa MCVV, Sena KXFR. *et al.* Study of the *Enterobacteriaceae* Group CESP (*Citrobacter*, *Enterobacter*, *Serratia*, *Providencia*, *Morganella* and *Hafnia*): A Review. In: *The Battle Against Microbial Pathogens: Basic*

- Science, Technological Advances and Educational Programs. 2015; Publisher: Formatex 2015.
32. World Health Organization. WHO. Antimicrobial Resistance in the WHO African Region: a systematic literature review. *Antimicrobial resistance in the WHO African Region: a systematic literature review*. 2021.
 33. Sekwadi P, Smith AM, Maruma W, Mongologa G, Tsele G, Ngomane M, Tau N. *et al.* A Prolonged Outbreak of Enteric Fever Associated With Illegal Miners in the City of Matlosana, South Africa, November 2020–September 2022, *Open Forum. Inf. Dis.* 2024;11(3):ofae118, <https://doi.org/10.1093/ofid/ofae118>
 34. Fatimawali Kepel BJ, Gani MA, Tallei TE. Comparison of Bacterial Community Structure and Diversity in Traditional Gold Mining Waste Disposal Site and Rice Field by Using a Metabarcoding Approach. *Int. J. Microbiol.* 2020;7:1858732. <https://doi:10.1155/2020/1858732>.
 35. Cardona GI, Escobar MC, Acosta-González A, Marín P, Marqués S. Highly mercury-resistant strains from different Colombian Amazon ecosystems affected by artisanal gold mining activities. *Appl. Microbiol. Biotechnol.* 2022;106:2775–2793 <https://doi.org/10.1007/s00253-022-11860-y>
 36. Ojo GJ, Onile OS, Momoh AO, Oyeyemi BF, Omoboyede V, Fadahunsi AI, Onile T. Physiochemical analyses and molecular characterization of heavy metal-resistant bacteria from Ilesha gold mining sites in Nigeria. *J. Genetic Eng. Biotechnol.* 2023;21:172 <https://doi.org/10.1186/s43141-023-00607-5>
 37. Yan C, Wang F, Liu H, Liu H, Pu S, Lin F, Geng H, Ma S, Zhang Y, Tian Z, Chen H, Zhou B, Yuan R. Deciphering the toxic effects of metals in gold mining area: Microbial community tolerance mechanism and change of antibiotic resistance genes. *Environ. Res.* 2020;189:109869. <https://doi:10.1016/j.envres.2020.109869>
 38. Evans J, Hanoodi M, Wittler M. (2024). Amoxicillin Clavulanate In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024, Available from: <https://www.ncbi.nlm.nih.gov/books/NBK538164/> (Accessed August, 2024).
 39. Sethuvel DPM, Bakthavatchalam YD, Karthik M, Irulappan M, Shrivastava R, Periasamy H. *et al.* β -Lactam Resistance in ESKAPE Pathogens Mediated Through Modifications

- in Penicillin-Binding Proteins: An Overview. *Infect.Dis.Ther.*2023;12: 829–841. <https://doi.org/10.1007/s40121-023-00771-8>
40. Bui T, Patel P, Preuss CV. Cephalosporins. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan–. PMID: 31855361. [Cephalosporins - PubMed \(nih.gov\)](#) (Accessed August, 2024).
41. Jiang H, Yu T, Yang Y, Yu S, Wu J, Lin R, Li Y, Fang J, Zhu C. Co-occurrence of Antibiotic and Heavy Metal Resistance and Sequence Type Diversity of *Vibrio parahaemolyticus* Isolated from *Penaeus vannamei* at Freshwater Farms, Seawater Farms, and Markets in Zhejiang Province, China. *Front. Microbiol. Sec. Food Microbiol.* 2020;11:2020 <https://doi.org/10.3389/fmicb.2020.01294>
42. Edet UO, Bassey IU, Joseph AP. Heavy metal co-resistance with antibiotics amongst bacteria isolates from an open dumpsite soil. *Heliyon.* 2023;9:e13456.
43. Nosalova L, Willner J, Fornalczyk A. *et al.* Diversity, heavy metals, and antibiotic resistance in culturable heterotrophic bacteria isolated from former lead–silver–zinc mine heap in Tarnowskie Gory (Silesia, Poland). *Arch.Microbiol.* 2023;205:26 <https://doi.org/10.1007/s00203-022-03369-9>
44. Adebisi YA, Jimoh ND, Ogunkola IO, Uwizeyimana T, Olayemi AH. *et al.* The use of antibiotics in COVID-19 management: a rapid review of national treatment guidelines in 10 African countries. *Trop MedHealth.*2021;49:51 <https://doi.org/10.1186/s41182-021-00344-w>
45. Zanichelli V, Sharland M, Cappello B, Moja L, Getahun H, Pessoa-Silva C, Sati H, van Weezenbeek C, Balkhy H, Simão M, Gandra S, Huttner B. *The WHO AWaRe (Access, Watch, Reserve) antibiotic book and prevention of antimicrobial resistance.* *Bull World Health Organ.* 2023;101(4):290–6. <https://doi:10.2471/BLT.22.288614>