

The potential effects of Azolla as a growth promoter for *Labeo rohita* fingerlings

Abstract

The 60 days experiment was conducted for the introduction of Azolla into the diet of *Labeo Rohita* fingerlings resulted in significant variations ($p < 0.05$) in several growth parameters, including percentage weight gain, percentage length gain, specific growth rate (SGR), feed conversion ratio (FCR), and survival rate. While no significant difference in FCR was observed among treatments during the initial 0 to 45-day interval, a notable disparity emerged between 45 to 60 days, with the highest FCR recorded in T3 and the lowest in T1. Similarly, although there were no significant differences in SGR among treatments at various intervals, the highest SGR was observed in T3, while the lowest was in T1. Furthermore, percentages of weight increment were significantly different between the intervals of 15-30 days and 30-45 days, with the highest percentage gain observed in T3 and the lowest in T1. Likewise, percentages of length increment showed significant differences between the intervals of 15-30 days, with T1 exhibiting the highest percentage gain and T1 the lowest. The survival rate % of different treatments at the end of the experiment varied, with T3 showing the highest survival rate and T0 the lowest. Statistical analysis confirmed significant differences in percent length gain, percent weight gain, SGR, and FCR among the treatments. These findings underscore the complex interplay between fish species, feed nutrients, additives, and rearing conditions on the growth process of *Labeo Rohita* fingerlings. The observed variations in growth parameters highlight the potential of Azolla as a growth promoter in aquaculture practices. Further research is warranted to elucidate the underlying mechanisms driving these effects and to optimize Azolla supplementation protocols for enhanced growth performance and sustainability in *Labeo Rohita* aquaculture.

Introduction

Aquaculture is a promising field that can help to meet the food requirements of the growing global population. However, it is crucial to ensure that aquaculture practices are sustainable. In 2020, the production of aquaculture worldwide reached an all-time high of 122.6 million tonnes, with a total value of \$1.5 billion. Aquatic animals made up 87.5 million tonnes of this production, while algae accounted for 35.1 million tonnes. Aquatic plants are essential to the growth and body composition of fish, and they provide vital nutrients for them. In some cases, they can even replace up to 50% of commercial feeds. It is recommended to use locally grown aquatic plants because the chemical composition of these plants is significantly impacted by the environment in which they grow. Azolla is an aquatic plant with a high percentage of nutrients, including minerals, chlorophyll, carotenoids, amino acids and vitamins. As a result, it is a useful ingredient for fish ponds and can be used directly or indirectly. Azolla helps macrophytophagous fish gain weight directly and can be fed to fish in Azolla fish pond cultures. Furthermore, Azolla increases the amount of fish feces produced, which is then used as an organic (nitrogenous) fertilizer by bottom dwellers. This leads to an increase in the overall productivity of ponds. Additionally, Azolla's rapid decomposition rates

التعليق [M1]: Abstract should begin with a sentence regarding the importance and necessity of the research.

التعليق [M2]: Keywords:

التعليق [M3]: Source

التعليق [M4]: ??

make it an excellent substrate for microbial processing, such as composting, or for enhancing the detrital food chain before it is applied to ponds. Rohu is a preferred Indian major carp species for cultivation. Various supplementary feeds have been experimented with to increase fish growth and production per unit area. Azolla has been explored as a potential feed supplement, and while it shows promise, it may not be sufficient for high productivity when used alone. Therefore, it can serve as a beneficial supplement to natural feed in low-input aquaculture systems, potentially reducing reliance on fish meal and fish oil sourced from nature. Azolla meal has been found to be significant in the diets of Tilapia species (*Oreochromis niloticus*, *Tilapia mossambica*, *Tilapia zillii*) and members of the Cyprinidae family (*Labeo rohita*, *Catla catla*, *Labeo calbasu*, *Labeo fimbriatus*, *Ctenopharyngodon idella*, *Barbonymus gonionotus*), as observed in the reviewed literature.

التعليق [M5]: !!!!!

التعليق [M6]: Source

Material and Method

Experimental Fish:- The present study selected healthy rohu (*Labio rohita*) fingerlings from the fish hatchery of College of Fishery Science, NDVSU, Jabalpur (M.P.).

Experimental diets:-

Experimental diets was formulated comprising mustard oilcake, rice bran, soya-bean meal, wheat flour, carboxymethylcellulose (CMC) and mineral mixture. To create experimental diets, Azolla was replaced with other protein sources into the basal diet at various proportions: 0.0 g/kg (T₀ control), 200 g/kg (T₁), 300 g/kg (T₂), 400 g/kg (T₃). Approximately 27% protein-containing diet was prepared for the experiment. The diet preparation is given in Table No. 1

Table No. 1: The ingredients used for the preparation of different experimental diets (g/Kg)

S. No.	Ingredients	T ₀	T ₁	T ₂	T ₃
1.	Azolla	0	200	300	400
2.	Soya-bean meal	350	300	250	200
3.	Mustard oilcake	190	170	190	220
4.	Wheat flour	360	230	170	90
5.	Carboxymethylcellulose	40	40	30	30

Experimental design:- The experiment lasted for 60 days and was conducted at the Aquaculture Research Laboratory of the College of Fishery Science, NDVSU, Jabalpur. In each tank of 500 liters, five fingerlings of *Labeo rohita* were randomly distributed and triplicated. The stocked fingerlings appeared to be healthy and were fed 5% of their body weight per day. The growth and growth parameters were monitored at 15-day intervals.

Growth parameters:-

Several growth parameters were calculated to analyze the effect of various dietary sources. Initial & final length, initial & final weight, final mean body length, final mean body weight were measured. Specific growth rates (SGR), the percentage increase in length and weight gain, survival rate, feed conversion ratio (FCR) were calculated using following mathematical formulae.

Specific growth rate (SGR) = $\frac{\log \text{ final body weight} - \log \text{ initial body weight}}{\text{time}} \times 100$

$$\text{Percentage increment in length} = \frac{\text{Mean final length of fish} - \text{Mean initial length of fish (cm)} \times 100}{\text{Mean initial length of fish}}$$

$$\text{Percentage increment in weight} = \frac{\text{Mean final weight of fish} - \text{Mean initial weight of fish (g)} \times 100}{\text{Mean initial weight of fish}}$$

$$\text{Survival rate} = \frac{\text{Total number of fishes harvested}}{\text{Total number of stocked}} \times 100$$

$$\text{Feed Conversion Ratio (FCR)} = \frac{\text{Feed given}}{\text{Weight gain}}$$

Result & Discussion

Table No. 2: The feed conversion ratio for various treatments was presented at different time intervals in days.

Treatment	0-15 Days	15-30 Days	30-45 Days	45-60 Days
T ₀	2.32± 0.26	2.79± 0.45	2.78± 0.35	1.56 ^a ± 0.19
T ₁	1.88± 0.50	1.90± 0.03	3.23± 0.80	1.51 ^{ab} ± 0.14
T ₂	1.37±0.10	2.97± 0.81	1.72± 0.31	1.70 ^{ab} ± 0.39
T ₃	1.24±0.08	1.42±0.12	3.26± 0.75	2.86 ^c ± 0.13

Table No. 3: Specific growth rate for various treatments was presented at different time intervals in days.

Treatment	0-15 Days	15-30 Days	30-45 Days	45-60 Days
T ₀	0.78± 0.15	0.66± 0.10	0.64± 0.08	0.66± 0.05
T ₁	1.06± 0.31	0.94± 0.24	0.57± 0.01	0.57± 0.02
T ₂	1.23± 0.23	0.74± 0.30	1.14± 0.43	0.74± 0.13
T ₃	1.39± 0.37	1.19± 0.19	0.67± 0.28	1.07± 0.58

Table No. 4: Percentage increment in weight for various treatments was presented at different time intervals in days.

Treatment	0-15 Days	15-30 Days	30-45 Days	45-60 Days
T ₀	19.99± 2.59	40.71 ^a ± 6.82	64.27 ^a ± 10.04	77.07± 7.51
T ₁	27.59± 7.04	57.77 ^{ab} ± 8.24	82.16 ^{ab} ± 10.42	86.37± 8.51
T ₂	33.05± 2.46	55.78 ^{ab} ± 2.45	99.14 ^{abc} ± 9.83	92.71± 7.65
T ₃	36.49± 2.34	80.34 ^c ± 4.01	107.54 ^{abc} ± 1.20	76.08± 2.21

Table No. 5: Percentage increment in length for various treatments was presented at different time intervals in days.

Treatment	0-15 Days	15-30 Days	30-45 Days	45-60 Days
T ₀	6.79± 1.58	3.89 ^a ± 1.03	8.83± 0.44	17.54± 1.67
T ₁	9.48± 1.79	7.62 ^{ab} ± 0.46	7.15± 1.39	16.97± 1.27
T ₂	10.39± 1.43	9.40 ^b ± 0.60	6.01± 0.48	13.77± 0.52
T ₃	11.71± 1.10	11.95 ^c ± 1.85	7.23± 0.45	15.07± 0.39




Fig. 1: Survival rate (%) of different treatments at the end of the experiment

In the current investigation, the introduction of Azolla in the diet led to significant variations ($p < 0.05$) in parameters such as percentage weight gain, percentage length gain, specific growth rate, feed conversion ratio, and survival rate, highlighting the intricate influence of fish species, feed nutrients, additives, and rearing conditions on the growth process. There is no significant difference in FCR observed in all treatments from 0 to 45 days intervals but between 45-60 days a significant difference was observed between the treatments. The highest value of FCR was found in T_3 (3.26 ± 0.75) and the lowest was also found in T_3 (1.24 ± 0.08) (Table no. 2). There is no significant difference was observed in SGR between all treatments at different day intervals. The highest value of SGR was found in T_3 (1.39 ± 0.37) and the lowest value was found in T_1 (0.57 ± 0.01) respectively (Table no. 3).

Percentages of weight increment were found to be significantly different between the intervals of 15-30 days and 30-45 days. Table no. 4 indicates that the highest percentage gain in weight was observed in T_3 (107.54 ± 1.20), whereas the lowest value was found in T_1 (19.99 ± 2.59). Similarly, percentages of length increment were found to be significantly different between the intervals of 15-30 days. Table no. 5 shows that the highest percentage gain in length was observed in T_1 (17.54 ± 1.67), whereas the lowest value was also found in T_1 (3.89 ± 1.03). Fig. 1 illustrates the survival rate % of different treatments at the end of the experiment. It is noteworthy that the highest survival rate was observed in T_3 , whereas the lowest was found in T_0 . The statistical analysis reveals significant differences in percent length gain, percent weight gain, specific growth rate (SGR), and feed conversion ratio (FCR). The results of present investigation were also found in several research. In the study conducted by Kumari et al. (2017), it was observed that dietary supplementation of Azolla had a beneficial impact on fish growth. Compared to the control group, the treatments

showed significant enhancements in weight gain, percent weight gain, specific growth rate (SGR), food conversion ratio (FCR), and gross conversion efficiency (GCE). The researchers also suggested that substituting dietary components with up to 20% Azolla could be a viable feeding strategy for monosex Nile tilapia. This approach has the potential to enhance fish growth, quality, profitability, and sustainability within the tilapia production industry (Refaey *et al.*, 2023).

Conclusion

The findings of the current study indicates the significant role of azolla as a growth promoter for aquatic weed when incorporated into the basal diet consisting of groundnut oil cake, rice bran, soya bean meal, wheat flour, and mineral mixture for *Labeo rohita* fingerlings. The results indicate improved percentage weight gain, percentage length gain, specific growth rate, feed conversion ratio, and survival rate in fingerlings fed with Azolla-supplemented diets. In summary, the study concludes that Azolla, at a maximum level of 300g/kg and 400g/kg (in dried pellet form), serves as a suitable dietary protein supplement for *Labeo rohita*, without any detrimental effects on growth performance, survival rate, and economic parameters. By enhancing growth rates, improving feed conversion efficiency, and maintaining economic feasibility, Azolla supplementation offers promising prospects for enhancing the productivity and profitability of *Labeo rohita* aquaculture. Further research and practical trials are recommended to validate these findings and explore the broader implications of integrating azolla into aquafeed formulations for commercial fish production.

References

1. Abo El-Ward, A., Eid, A. E., Mohamed, K. A., Abd-elfattah, B., & Hasan, M. A. (2016). Growth performance of Nile tilapia (*Oreochromis niloticus*) fingerlings fed diet supplemented with different of *Spirulina platensis* levels. *Egyptian Journal of Animal Production*, 53(3), 181-190.
2. Ayyappan, S., & Jena, J. K. (2003). Grow-out production of carps in India. *Journal of Applied Aquaculture*, 13(3-4), 251-282.
3. Debnath, D., Yengkokpam, S., Bhattacharjya, B. K., Biswas, P., Prakash, C., Kohli, M. P. S., & Sharma, A. P. (2018, March). Effect of dietary incorporation of dry-powdered

التعليق [M7]: References citations and use in the article text MUST be in Chronological order
The References list must be in Alphabetical order.
Cross-Check the References...
There should be NO capital letters inside the Title of the article in the Reference.
All the names must be in italics, please.

water hyacinth (*Eichhornia crassipes*) meal on growth and digestibility of *Labeo rohita* fingerlings. In *Proceedings of the Zoological Society*, 71, 74-82).

4. Ebrahim, M. S. M., Zeinhom, M. M., & Abou-Seif, R. A. (2007). Response of Nile tilapia (*Oreochromis niloticus*) fingerlings to diets containing Azolla meal as a source of protein. *Journal of the Arabian*, 2(1), 54-69.
5. Gangadhar, B., Sridhar, N., Saurabh, S., Raghavendra, C. H., Hemaprasanth, K. P., Raghunath, M. R., & Jayasankar, P. (2015). Effect of Azolla-incorporated diets on the growth and survival of *Labeo fimbriatus* during fry-to-fingerling rearing. *Cogent Food & Agriculture*, 1(1), 1055539.
6. Gangadhar, B., Sridhar, N., Saurabh, S., Raghavendra, C. H., Hemaprasanth, K. P., Raghunath, M. R., & Jayasankar, P. (2014). Growth Response of *Cirrhinus mrigala* fry to Azolla (*Azolla pinnata*)-incorporated Diets. *Fishery Technology* 51 (2014),156 – 161
7. Hassan, M. S., & Edwards, P. (1992). Evaluation of duckweed (*Lemna perpusilla* and *Spirodela polyrrhiza*) as feed for Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 104(3-4), 315-326.
8. Ibrahim, A., Abouelella, S., Adam, H., Hassan, N., & Abdallah, R. (2021). Effect of *Azolla pinnata* on growth performance and survival rate of fingerlings of grass carp fish *Ctenopharyngodon idellus* (Valenciennes, 1844). *Aswan University Journal of Environmental Studies*, 2(4), 249-258.
9. Kotnala, S., Dhar, P., Das, P., & Chatterji, A. (2010). Growth performance of fingerlings of the Indian major carp *Catla catla* (Ham.) fed with feeds supplemented with different seaweeds. *Pertanika Journal of Science. & Technology*, 18(2), 255-62.
10. Kumari, R., Ojha, M. L., Saini, V. P., & Sharma, S. K. (2017). Effect of Azolla supplementation on growth of rohu (*Labeo rohita*) fingerlings. *Journal of entomology and zoology studies*, 5(4), 1116-19.
11. Mosha, S. S. (2018). A review on significance of Azolla meal as a protein plant source in finfish culture. *Journal of Aquaculture Research and Development*, 9(7).
12. Refaey, M. M., Mehrim, A. I., Zenhom, O. A., Areeda, H. A., Ragaza, J. A., & Hassaan, M. S. (2023). Fresh Azolla, *Azolla pinnata* as a complementary feed for *Oreochromis niloticus*: growth, digestive enzymes, intestinal morphology, physiological responses, and flesh quality. *Aquaculture Nutrition*, 2023. enzymes, intestinal morphology, physiological responses, and flesh quality. *Aquaculture Nutrition*, 2023.

UNDER PEER REVIEW