

Fixed point theorem utilizing rational-type integral contraction for multivalued mappings in cone $b - metric - like$ space

Abstract: We prove common fixed point theorem for multivalued mappings in a cone $b - metric - like$ space by using a rational-type integral contraction condition. Some sufficient conditions are obtained for the existence of common fixed points of multivalued mappings defined on this space . The obtained results generalize and extend some known results for fixed point theorems in cone $b - metric - like$ space. Some corollaries are obtained as special cases of the main theorem. An example is also given to illustrate the applicability of the obtained results.

Key words: Fixed point, Cone $b - metric - like$ space, Integral contraction.

AMS Subject Classification (2020): 47H10, 54E50, 47H04.

1. INTRODUCTION:

Fixed point theorems in generalized cone metric spaces are equal to those in ordinary metric spaces, according to Du [1]. Specifically, the author demonstrated that the well-known fixed-point theorems of Banach, Kannan, and Chatterjea in topological vector spaces and cone metric may be readily obtained from the standard metric space setup through a straightforward modification, specifically the use of a scalarization function.

As a generalization of $b - metric - like$ spaces, Fernandez et al. [2] introduced cone $b - metric - like$ spaces over Banach algebra in 2017. They also studied the fixed point of generalized contractions and expansive mapping in the context of cone $b - metric - like$ spaces over Banach algebras with a nonnormal cone.

Reich [3] developed a novel kind of contraction in 1971 that we refer to as Re-

ich contraction. The Banach contraction and the Kannan contraction are the two prominent contractions that it generalizes. The Banach contraction principle was expanded by Nadler [4] from single-valued to multivalued contraction maps, taking into account the metric defined on bounded and closed subsets of a non-empty set Z . Furthermore, by introducing the concepts of a b -metric space [5], a b -metric-like space [6], a partial metric space [7], a quasi b -metric space [8 and 9], a cone rectangular metric space [10], a new extended b -metric space [11], cone metric spaces [12 and 13], cone b -metric spaces [14], and extended b -metric spaces [15], the fixed point theory is widely accepted by altering the broad structure of underlying space.

The purpose of this work is to extend fixed point theory to the more general setting of cone b -metric-like spaces. It develops new common fixed point results for multivalued mappings using rational-type contraction conditions. The results generalize and unify several existing theorems in the literature. Ultimately, the work provides a broader framework for solving nonlinear problems and example arising in analysis.

2. PRELIMINARIES:

In this section, we recall some basic concepts and results which will be used throughout the paper. These notions form the foundation of fixed point theory in generalized metric structures, particularly in cone b -metric-like spaces.

Let Y be a real Banach space and $V \subset Y$ be a subset.

Definition: 2.1. [16 and 17] A subset V of a real Banach space Y is called a *cone* if it satisfies the following conditions:

- (i) V is closed, nonempty and $V \neq \{\vartheta\}$;
- (ii) for all $c, d \geq 0$ and $z, x \in V$, we have $cz + dx \in V$;
- (iii) $V \cap (-V) = \{\vartheta\}$.

The cone V induces a partial ordering $\preceq$ on Y defined by

$$z \preceq x \iff x - z \in V.$$

We write $z \prec x$ if $z \preceq x$ and $z \neq x$, and $z \ll x$ if $x - z$ belongs to the interior of V .

Definition: 2.2. [16] A cone V is said to be *normal* if there exists a constant $L > 0$ such that

$$\vartheta \preceq z \preceq x \implies \|z\| \leq L\|x\|,$$

for all $z, x \in Y$. The least such L is called the normal constant of V .

The concept of cone metric spaces was later generalized to cone *b-metric* spaces, and further to cone *b-metric-like* spaces, which allow more flexibility in handling nonlinear problems.

Definition: 2.3. [18] Let Z be a nonempty set and $S \geq 1$ be a real number. A mapping $d : Z \times Z \rightarrow Y$ is called a *cone b-metric* if for all $z, x, w \in Z$, the following conditions hold:

- (i) $\vartheta \preceq d(z, x)$ and $d(z, x) = \vartheta \iff z = x$;
- (i) $d(z, x) = d(x, z)$;
- (i) $d(z, x) \preceq S(d(z, w) + d(w, x))$.

Definition: 2.4. [19] A mapping $d : Z \times Z \rightarrow Y$ is called a *cone b-metric-like* if it satisfies all the conditions of a cone *b-metric* except that $d(z, z)$ need not be equal to ϑ , that is,

$$d(z, z) \neq \vartheta \quad \text{in general.}$$

The pair (Z, d) is called a *cone b-metric-like space*.

This generalization plays an important role in extending fixed point results, since it allows self-distance to be non-zero, which appears naturally in various applications.

Definition: 2.5. [18] Let (Z, d) be a cone *b-metric-like* space. A sequence $\{z_p\}$ for all $p \in \mathbb{N}$ in Z is said to be:

(i) *Cauchy* if for every $c \in Y$ with $\vartheta \ll c$, there exists N such that for all $p, m \geq N$,

$$d(z_p, z_m) \ll c;$$

(i) *convergent* to $z \in Z$ if for every $c \in Y$ with $\vartheta \ll c$, there exists N such that for all $n \geq N$,

$$d(z_p, z) \ll c.$$

Definition: 2.6. [20] A cone *b-metric-like* space (Z, d) is said to be *complete* if every Cauchy sequence in Z converges to a point in Z .

In cone *b-metric-like* spaces, the concept of a fixed point differs slightly from the classical case.

Definition: 2.7. [21] Let $G : Z \rightarrow Z$ be a mapping. A point $z \in Z$ is called a *metric-like fixed point* of G if

$$d(z, Gz) = d(z, z).$$

For multivalued mappings, we use the following notion.

Definition: 2.8. [22] Let $G : Z \rightarrow \mathcal{S}_M(Z)$ be a multivalued mapping. A point $z \in Z$ is said to be a *fixed point* of G if

$$z \in Gz.$$

Throughout this paper, we denote by $\mathcal{S}_M(Z)$ the family of all nonempty subsets of Z . We shall consider mappings whose values are nonempty, closed and bounded subsets of Z .

We also make use of the following class of functions.

Definition: 2.9. [23] A function $\varphi : V \rightarrow [0, \infty)$ is said to belong to the class Φ if it satisfies:

- (i) φ is continuous and non-decreasing;
- (ii) $\varphi(w) = 0$ if and only if $w = \vartheta$.

These concepts will be used to establish our main results concerning common fixed points of multivalued mappings under integral-type rational contractions.

3. MAIN RESULT:

Theorem: 3.1. Let (Z, d) be a complete cone b -metric-like space with coefficient $S \geq 1$ over a real Banach space Y with cone V . Assume that:

- (i). $d(z, x) = \vartheta \Rightarrow z = x$ for all $z, x \in Z$
- (ii). $G, H : Z \rightarrow S_M(Z)$ are two multivalued mappings.
- (iii). There exists a function $\phi : S_M(Z) \rightarrow [0, \infty)$ which is continuous, non-decreasing and satisfies $\phi(w) = 0$ if and only if $w = \vartheta$, and a constant $k \in (0, \frac{1}{s})$ such that

$$\int_0^{d(Gz, Hx)} \phi(w) dw \preceq k \int_0^{d(z, x)} \phi(w) dw \quad \text{for all } z, x \in Z \quad (1)$$

- (iv). For each $z \in Z$, Gz and Hx are nonempty, closed and bounded subsets of Z .

Then G and H have a common fixed point, that is, there exists $q \in Z$ such that

$$q \in Gq \cap Hq.$$

Proof: Let $z_0 \in Z$ be arbitrary. Choose $z_1 \in Gz_0$, $z_2 \in Hz_1$, $z_3 \in Gz_2$, and so on. Thus, we construct a sequence $\{z_p\}$ in Z such that

$$z_{2p+1} \in Gz_{2p}, \quad z_{2p+2} \in Hz_{2p+1}, \quad \text{for all } p \in \mathbb{N}.$$

By the stated contractive condition, we have

$$\int_0^{d(z_{p+1}, z_{p+2})} \phi(w) dw \preceq k \int_0^{d(z_p, z_{p+1})} \phi(w) dw.$$

By the provided integral type contractive condition, for all $z, x \in Z$ we have

$$\int_0^{d(Gz, Hx)} \phi(w) dw \preceq k \int_0^{d(z, x)} \phi(w) dw. \quad (2)$$

Now, by using the construction of the sequence $\{z_p\}$ so that

$$z_{p+1} \in Gz_p, \quad z_{p+2} \in Hz_{p+1},$$

we obtain

$$d(z_{p+1}, z_{p+2}) \preceq d(Gz_p, Hz_{p+1}).$$

Given that ϕ is non-decreasing, it is evident that

$$\int_0^{d(z_{p+1}, z_{p+2})} \phi(w) dw \preceq \int_0^{d(Gz_p, Hz_{p+1})} \phi(w) dw.$$

By using the contractive condition, we obtain

$$\int_0^{d(z_{p+1}, z_{p+2})} \phi(w) dw \preceq k \int_0^{d(z_p, z_{p+1})} \phi(w) dw. \quad (3)$$

We now go over this inequality again. We have for $p = 0, p = n$

$$\int_0^{d(z_1, z_2)} \phi(w) dt \preceq k \int_0^{d(z_0, z_1)} \phi(w) dw.$$

For $p = 1$, we obtain

$$\int_0^{d(z_2, z_3)} \phi(w) dw \preceq k \int_0^{d(z_1, z_2)} \phi(w) dw \preceq k^2 \int_0^{d(z_0, z_1)} \phi(w) dw.$$

Proceeding inductively, assume that

$$\int_0^{d(z_p, z_{p+1})} \phi(w) dw \preceq k^p \int_0^{d(z_0, z_1)} \phi(w) dw.$$

Next, applying the contractive condition once more,

$$\int_0^{d(z_{p+1}, z_{p+2})} \phi(w) dw \preceq k \int_0^{d(z_p, z_{p+1})} \phi(w) dw \preceq k^{p+1} \int_0^{d(z_0, z_1)} \phi(w) dw.$$

Thus, by mathematical induction, for all $p \in \mathbb{N}$,

$$\int_0^{d(z_p, z_{p+1})} \phi(w) dw \preceq k^p \int_0^{d(z_0, z_1)} \phi(w) dw. \quad (4)$$

Since $0 < k < \frac{1}{r} < 1$, we have $k^p \rightarrow 0$ as $p \rightarrow \infty$. Hence,

$$\int_0^{d(z_p, z_{p+1})} \phi(w) dw \rightarrow \vartheta.$$

Using the property of ϕ , it follows that

$$d(z_p, z_{p+1}) \rightarrow \vartheta.$$

Now, using the cone b -metric-like inequality, for $m > n$ we have

$$d(z_p, z_m) \preceq r [d(z_p, z_{p+1}) + d(z_{p+1}, z_{p+2}) + \cdots + d(z_{m-1}, z_m)].$$

Using the above estimates and the fact that $k < 1$, we conclude that

$$d(z_p, z_m) \rightarrow \vartheta \quad \text{as } p, m \rightarrow \infty.$$

Thus, $\{z_p\}$ is a Cauchy sequence in Z .

Since (Z, d) is complete, there exists $u \in X$ such that

$$z_p \rightarrow q.$$

Next, we show that $q \in Hq$. From the contractive condition, we have

$$\int_0^{d(z_{2p+1}, Hq)} \phi(w) dw \preceq k \int_0^{d(z_{2p}, q)} \phi(w) dw.$$

Taking the limit as $p \rightarrow \infty$, the right-hand side tends to ϑ , hence

$$d(z_{2p+1}, Hq) \rightarrow \vartheta.$$

Since $z_{2p+1} \rightarrow q$ and Hq is closed, it follows that $q \in Hq$.

Similarly, we show that $q \in Gq$. Indeed,

$$\int_0^{d(z_{2p+2}, Gq)} \phi(w) dw \preceq k \int_0^{d(z_{2p+1}, q)} \phi(w) dw.$$

Taking the limit, we get

$$d(z_{2p+2}, Gq) \rightarrow \vartheta,$$

which implies that $q \in Gq$.

Therefore,

$$q \in Gq \cap SHq,$$

and hence q is a common fixed point of G and H .

Corollary: 3.2. Let (Z, d) be a complete cone b -metric-like space with coefficient $S \geq 1$ over a real Banach space Y with cone V . Assume that:

- (i). $d(z, x) = \vartheta \Rightarrow z = x$ for all $z, x \in Z$;
- (ii). $G : Z \rightarrow S_M(Z)$ is a multivalued mapping;
- (iii). There exists a function $\phi : P \rightarrow [0, \infty)$ which is continuous, non-decreasing and satisfies $\phi(w) = 0$ if and only if $w = \vartheta$, and a constant $k \in (0, \frac{1}{s})$ such

that

$$\int_0^{d(Gz, Gx)} \phi(w) dw \preceq k \int_0^{d(z, x)} \phi(w) dw \quad \text{for all } z, x \in Z \quad (5)$$

(iv). For each $z \in Z$, Gz is nonempty, closed and bounded.

Then G has a fixed point, that is, there exists $q \in Z$ such that

$$q \in Gq.$$

Proof: The result follows directly from the main theorem by taking $H = G$. Then all the conditions of the theorem are satisfied, and hence G has a fixed point in Z .

Corollary: 3.3. Let (Z, d) be a complete cone *b-metric-like* space with coefficient $S \geq 1$. Let $G : Z \rightarrow S_M(Z)$ be a multivalued mapping. Suppose there exists $k \in (0, \frac{1}{s})$ such that

$$d(Gz, Gx) \preceq k d(z, x) \quad \text{for all } z, x \in Z.$$

Then G has a fixed point in Z .

Corollary: 3.4. Let (Z, d) be a complete cone *b-metric-like* space with coefficient $S \geq 1$. Let $G, H : Z \rightarrow S_M(Z)$ be two multivalued mappings such that for some $k \in (0, \frac{1}{s})$,

$$\int_0^{d(Gz, Hx)} w dw \preceq k \int_0^{d(z, x)} w dw \quad \text{for all } z, x \in Z.$$

Then G and H have a common fixed point in Z .

Proof: Taking $\phi(w) = w$ in the main theorem, all conditions are satisfied. Hence, the result follows immediately.

Corollary: 3.5. Let (Z, d) be a complete cone *b-metric-like* space with coefficient $S \geq 1$. Let $G, H : Z \rightarrow S_M(Z)$ be two multivalued mappings such that there exists

$k \in (0, \frac{1}{s})$ satisfying

$$d(Gz, Hx) \preceq k d(z, x) \quad \text{for all } z, x \in Z.$$

Then G and H have a common fixed point in Z .

Proof: The result follows by choosing $\phi(w) = 1$ (or an appropriate constant function) in the integral type contraction and applying the main theorem.

Alternatively, the result can be obtained by considering a suitable constant function ϕ and reducing the integral inequality to the standard contraction form.

Example: 3.6. Let $Z = [0, 1]$ and let $\psi = C_{\mathbb{R}}^1[0, 1]$ be the set of all real-valued continuously differentiable functions on $[0, 1]$. Define the norm on ψ by

$$\|g\| = \|g\|_{\infty} + \|g'\|_{\infty}.$$

Let $V = \{g \in \psi : g(w) \geq 0 \text{ for all } w \in [0, 1]\}$ be a cone in ψ .

Define $d : Z \times Z \rightarrow \psi$ by

$$d(z, x)(w) = |z - x|^m e^w + 1, \quad m > 1.$$

Then (Z, d) is a cone b -metric-like space with coefficient $s = 2^m$, since for all $z, x, y \in Z$,

$$d(z, x)(t) \preceq r(d(z, x)(t) + d(x, y)(w)).$$

Note that $d(z, z)(w) = 1 \neq \vartheta$, hence (Z, d) is not a cone b -metric space, but it is a cone b -metric-like space.

Now define two multivalued mappings $G, H : Z \rightarrow PQ(Z)$ by

$$Gz = \left[0, \frac{z}{4}\right], \quad Hz = \left[0, \frac{z}{6}\right].$$

Clearly, for each $z \in Z$, Gz and Hx are non-empty, closed and bounded subsets of Z .

Now, for $z, x \in Z$ with $z \geq x$, we estimate

$$d(Gz, Hx) \preceq \left| \frac{z}{4} - \frac{x}{6} \right|^p e^w + 1.$$

Since

$$\left| \frac{z}{4} - \frac{x}{6} \right| \leq \frac{1}{4} |z - x|,$$

we obtain

$$d(Gz, Hx) \preceq \frac{1}{4^m} |z - x|^m e^w + 1.$$

Now, choose $\phi(w) = w$ (linear function). Then

$$\int_0^{d(Gz, Hx)} \phi(w) dw \preceq \frac{1}{4^m} \int_0^{d(z, x)} \phi(w) dw.$$

Let $k = \frac{1}{4^m}$. Since $m > 1$, we have $k < \frac{1}{r}$.

Thus, all the conditions of the main theorem are satisfied. Therefore, G and H have a common fixed point in Z .

In fact, $q = 0$ is a common fixed point since

$$0 \in G0 \cap H0.$$

References

- [1] Du, W. S., and Karapinar, E. (2013). A note on cone b -metric and its related results: Generalizations or equivalence. *Fixed Point Theory and Applications*, 2013(1), Article ID. 210.
- [2] Fernandez, J., Malviya, N., and Shukla, S. (2017). Cone b-metric-like spaces over Banach algebra and fixed point theorems with the application. *Asian Journal of Mathematics and Computer Research*, 18(2), 49–66.
- [3] Reich, S. (1971). Some remarks concerning contraction mappings. *Canadian Mathematical Bulletin*, 14(1), 121–124.
- [4] Nadler, S. B. (1969). Multi-valued contraction mappings. *Pacific Journal of Mathematics*, 30(2), 475–488.
- [5] Bakhtin, I. A. (1989). The contraction mapping principle in quasi metric spaces. *Functional Analysis, Gosudarstv. Ped. Inst. Unianowsk*, 30, 26–37.
- [6] Zada, B., Riaz, U. (2016). Some fixed point results on multiplicative b -metric-like spaces. *Turkish Journal of Analysis and Number Theory*, 4(5), 118–131.
- [7] Aydi, H. (2011). Some coupled fixed point results on partial metric spaces. *International Journal of Mathematics and Mathematical Sciences*, 2011, 11, Article ID. 647091.
- [8] Klin-eam, C., and Suanoom, C. (2015). Dislocated $quasi - b - metric$ spaces and fixed point theorems for cyclic contractions. *Fixed Point Theory and Applications*, 2015(1), 12, Article ID. 74.
- [9] Rahman, M. U., and Sarwar, M. (2016). Dislocated quasi b-metric space and fixed point theorems. *Electronic Journal of Mathematical Analysis and Applications*, 4(2), 16–24.

- [10] Jleli, M., and Samet, B. (2009). The Kannan's fixed point theorem in a cone rectangular metric space. *Journal of Nonlinear Science and Applications*, 2(3), 161–167.
- [11] Samreen, M., Kamran, T., and Postolache, M. (2018). Extended b – metric space, extended b – comparison function and nonlinear contractions. *University Politehnica of Bucharest Scientific Bulletin Series A*. 80(4), 21–28.
- [12] Rezapour, S., and Hamlbarani, R. (2008). Some notes on the paper Cone metric spaces and fixed point theorems of contractive mappings. *Journal of Mathematical Analysis and Applications*, 345(2), 719–724.
- [13] Kadelburg, Z., and Radenović, S. (2013). A note on various types of cones and fixed point results in cone metric spaces. *Asian Journal of Mathematics and Applications*, 2013, 7, Article ID. ama0104.
- [14] Kutbi, M. A., Ahmad, J., Al-Mazrooei, A. E., and Hussain, N. (2018). Multi-valued fixed point theorems in cone b – metric spaces over Banach algebra with applications. *Journal of Mathematics and Analysis*, 9(1), 52–64.
- [15] Shatanawi, W., Abodayeh, K., and Mukheimer, A. (2018). Some fixed point theorems in extended b – metric spaces. *University Politehnica of Bucharest Scientific Bulletin Series A*. 80(4), 71–78.
- [16] Huang, L. G., & Zhang, X. (2007). Cone metric spaces and fixed point theorems of contractive mappings. *Journal of Mathematical Analysis and Applications*, 332(2), 1468–1476.
- [17] Rezapour, S., & Hamlbarani, R. (2008). Some notes on the paper Cone metric spaces and fixed point theorems of contractive mappings. *Journal of Mathematical Analysis and Applications*, 345(2), 719–724.

- [18] Hussain, N., & Shah, M. H. (2011). KKM mappings in cone b-metric spaces. *Computers & Mathematics with Applications*, 62(4), 1677–1684.
- [19] Fernandez, J., Malviya, N., & Shukla, S. (2017). Cone b-metric-like spaces over Banach algebra and fixed point theorems with applications. *Asian Journal of Mathematics and Computer Research*, 18(2), 49–66.
- [20] Alghamdi, M. A., Hussain, N., & Salimi, P. (2013). Fixed point and coupled fixed point theorems on b-metric-like spaces. *Journal of Inequalities and Applications*, 2013(1), 25 Article ID. 402.
- [21] Amini-Harandi, A. (2012). Metric-like spaces, partial metric spaces and fixed points. *Fixed Point Theory and Applications*, 2012(1), 11, Article ID. 204.
- [22] Chen, C. F., Dong, J., & Zhu, C. X. (2015). Some fixed point theorems in b-metric-like spaces. *Fixed Point Theory and Applications*, 2015(1), 13, Article ID. 122.
- [23] Aydi, H., Sahmim, S., & Mehmood, N. (2018). Common fixed point theorems for $(T, g)_F$ -contraction in b-metric-like spaces. *Journal of Inequalities and Applications*, 2018(1), 19 Article ID. 119.